



Europäisches Patentamt  
European Patent Office  
Offic européen des brevets



Publication number: **0 599 274 A1**

**EUROPEAN PATENT APPLICATION**

Application number: **93118837.9**

Date of filing: **23.11.93**

Int. Cl.<sup>5</sup>: **C12N 15/12, C07K 13/00,  
C12P 21/08, C12N 5/10,  
G01N 33/68, G01N 33/577,  
A61K 39/395**

Priority: **24.11.92 US 981165**

Date of publication of application:  
**01.06.94 Bulletin 94/22**

Designated Contracting States:  
**AT BE CH DE DK ES FR GB GR IE IT LI LU MC  
NL PT SE**

Applicant: **Bristol-Myers Squibb Company**  
**345 Park Avenue**  
**New York, N.Y. 10154(US)**

Inventor: **Plowman, Gregory D.**  
**1000 Union Street No. C.**

**Seattle, WA 98101(US)**  
Inventor: **Culouscou, Jean-Michel**  
**3034 W. Viewmont Way W.**  
**Seattle, WA 98199(US)**  
Inventor: **Shoyab, Mohammed**  
**2405 Westmont Way W.**  
**Seattle, WA 98199(US)**


Representative: **Kinzebach, Werner, Dr. et al**  
**Patentanwälte**  
**Reitstötter, Kinzebach und Partner**  
**Postfach 86 06 49**  
**D-81633 München (DE)**

**HER4, a human receptor tyrosine kinase of the epidermal growth factor receptor family.**

The molecular cloning, expression, and biological characteristics of a novel receptor tyrosine kinase related to the epidermal growth factor receptor, termed HER4/p180<sup>erbB4</sup>, are described. A HER4 ligand capable of inducing cellular differentiation of breast cancer cells is also disclosed. In view of the expression of HER4 in several human cancers and in certain tissues of neuronal and muscular origin, various diagnostic and therapeutic uses of HER4-derived and HER4-related biological compositions are provided.

**EP 0 599 274 A1**

FIGURE 5



## 1. INTRODUCTION

The present invention is generally directed to a novel receptor tyrosine kinase related to the epidermal growth factor receptor, termed HER4/p180<sup>erbB4</sup> ("HER4"), and to novel diagnostic and therapeutic compositions comprising HER4-derived or HER4-related biological components. The invention is based in part upon applicants' discovery of human HER4, its complete nucleotide coding sequence, and functional properties of the HER4 receptor protein. More specifically, the invention is directed to HER4 biologics comprising, for example, polynucleotide molecules encoding HER4, HER4 polypeptides, anti-HER4 antibodies which recognize epitopes of HER4 polypeptides, ligands which interact with HER4, and diagnostic and therapeutic compositions and methods based fundamentally upon such molecules. In view of the expression of HER4 in several human cancers and in certain tissues of neuronal and muscular origin, the present invention provides a framework upon which effective biological therapies may be designed. The invention is hereinafter described in detail, in part by way of experimental examples specifically illustrating various aspects of the invention and particular embodiments thereof.

## 2. BACKGROUND OF THE INVENTION

Cells of virtually all tissue types express transmembrane receptor molecules with intrinsic tyrosine kinase activity through which various growth and differentiation factors mediate a range of biological effects (reviewed in Aaronson, 1991, *Science* 254: 1146-52). Included in this group of receptor tyrosine kinases (RTKs) are the receptors for polypeptide growth factors such as epidermal growth factor (EGF), insulin, platelet-derived growth factor (PDGF), neurotrophins (i.e., NGF), and fibroblast growth factor (FGF). Recently, the ligands for several previously characterized receptors have been identified, including ligands for c-kit (steel factor), met (hepatocyte growth factor), trk (nerve growth factor) (see, respectively, Zsebo et al., 1990, *Cell* 63: 195-201; Bottardo et al., 1991, *Science* 251: 802-04; Kaplan et al., 1991, *Nature* 350: 158-160). In addition, the soluble factor NDF, or heregulin- $\alpha$  (HRG- $\alpha$ ), has been identified as the ligand for HER2, a receptor which is highly related to HER4 (Wen et al., 1992, *Cell* 69:559-72; Holmes et al., 1992 *Science* 256:1205-10). However, at present, the ligands for a number of isolated and/or characterized receptor tyrosine kinases have still not been identified, including those for the eph, eck, elk, ret, and HER3 receptors.

Biological relationships between various human malignancies and genetic aberrations in growth factor-receptor tyrosine kinase signal pathways are known to exist. Among the most notable such relationships involve the EGF receptor (EGFR) family of receptor tyrosine kinases (see Aaronson, *supra*). Three human EGFR-family members have been identified and are known to those skilled in the art: EGFR, HER2/p185<sup>erbB2</sup>, and HER3/p160<sup>erbB3</sup> (see, respectively, Ullrich et al., 1984, *Nature* 309: 418-25; Coussens et al., 1985, *Science* 230: 1132-39; and Plowman et al., 1990, *Proc. Natl. Acad. Sci. U.S.A.* 87: 4905-09). EGFR-related molecules from other species have also been identified.

The complete nucleotide coding sequence of other EGFR-family members has also been determined from other organisms including: the drosophila EGFR ("DER": Livneh, E. et al., 1985, *Cell* 40: 599-607), nematode EGFR ("let-23": Aroian, R.V. et al., 1990, *Nature* 348: 693-698), chicken EGFR ("CER": Lax, I. et al., 1988, *Mol. Cell. Biol.* 8: 1970-1978), rat EGFR (Petch, L.A. et al., 1990, *Mol. Cell. Biol.* 10: 2973-2982), rat HER2/neu (Bargmann, C.I. et al., 1986, *Nature*, 319: 226-230) and a novel member isolated from the fish and termed *Xiphophorus* melanoma related kinase ("Xmrk": Wittbrodt, J. et al., 1989, *Nature* 342: 415-421). In addition, PCR technology has led to the isolation of other short DNA fragments that may encode novel receptors or may represent species-specific homologs of known receptors. One recent example is the isolation tyro-2 (Lai, C. and Lemke, G., 1991, *Neuron* 6: 691-704) a fragment encoding 54 amino acids that is most related to the EGFR family.

Overexpression of EGFR-family receptors is frequently observed in a variety of aggressive human epithelial carcinomas. In particular, increased expression of EGFR is associated with more aggressive carcinomas of the breast, bladder, lung and stomach (see, for example, Neal et al., 1985, *Lancet* 1: 366-68; Sainsbury et al., 1987, *Lancet* 1: 1398-1402; Yasui et al., 1988, *Int. J. Cancer* 41: 211-17; Veale et al., 1987, *Cancer* 55: 513-16). In addition, amplification and overexpression of HER2 has been associated with a wide variety of human malignancies, particularly breast and ovarian carcinomas, for which a strong correlation between HER2 overexpression and poor clinical prognosis and/or increased relapse probability have been established (see, for example, Slamon et al., 1987, *Science* 235: 177-82, and 1989, *Science* 244: 707-12). Overexpression of HER2 has also been correlated with other human carcinomas, including carcinoma of the stomach, endometrium, salivary gland, bladder, and lung (Yokota et al., 1986, *Lancet* 1: 765-67; Fukushigi et al., 1986, *Mol. Cell. Biol.* 6: 955-58; Yonemura et al., 1991, *Cancer Res.* 51: 1034; Weiner et al., 1990, *Cancer Res.* 50: 421-25; Geurin et al., 1988, *Oncogene Res.* 3:21-31; Semba et al., 1985, *Proc. Natl. Acad.*

Sci. U.S.A. 82: 6487-6501; Zhau et al., 1990, Mol. Carcinog. 3: 354-57; McCann et al., 1980, Cancer 65: 88-92). Most recently, a potential link between HER2 overexpression and gastric carcinoma has been reported (Jeahne et al., 1992, J. Cancer Res. Clin. Oncol. 118: 474-79). Finally, amplified expression of the recently described HER3 receptor has been observed in a wide variety of human adenocarcinomas (Poller et al., 1992, J. Path., in press; Krause et al., 1989, Proc. Natl. Acad. Sci. U.S.A. 86: 8183-87; European Patent Application No. 91301737, published 9.4.91, EP 444 861).

Several structurally related soluble polypeptides capable of specifically binding to EGFR have been identified and characterized, including EGF, transforming growth factor- $\alpha$  (TGF- $\alpha$ ), amphiregulin (AR), heparin-binding EGF (HB-EGF), and vaccinia virus growth factor (VGF) (see, respectively, Savage et al., 1972, J. Biol. Chem. 247: 7812-21; Marquardt et al., 1984, Science 223: 1078-82; Shoyab et al., 1989, Science 243: 1074-76; Higashiyama et al., 1991, Science 251: 938-39; Twardzik et al., 1985, Proc. Natl. Acad. Sci. U.S.A. 82: 5300-04). Despite the close structural relationships among receptors of the EGFR-family, none of these ligands has been conclusively shown to interact with HER2 or HER3.

Recently, several groups have reported the identification of specific ligands for HER2. Some of these ligands, such as gp30 (Lupu et al., 1990, Science 249: 1552-55; Bacus et al., 1992, Cell Growth and Differentiation 3: 401-11) interact with both EGFR and HER2, while others are reported to bind specifically to HER2 (Wen et al., 1992, Cell 69: 559-72; Peles et al., 1992, Cell 69: 205-16; Holmes et al., 1992, Science 258: 1205-10; Lupu et al., 1992, Proc. Natl. Acad. Sci. U.S.A. 89: 2287-91; Huang et al., 1992, J. Biol. Chem. 276: 11508-121). The best characterized of these ligands are neu differentiation factor (NDF) purified and cloned from ras-transformed Rat1-EJ cells (Wen et al., Peles et al., *supra*), and the heregulins (HRF- $\alpha$ , - $\beta$ 1, - $\beta$ 2, - $\beta$ 3), purified and cloned from human MDA-MB-231 cells (Holmes et al., *supra*). NDF and HRF- $\alpha$  share 83% sequence identity and appear to be the rat and human homologs of the same protein. Both of these proteins are similar size (44-45 kDa), increase tyrosine phosphorylation of HER2 in MDA-MB-453 cells and not the EGF-receptor, and have been reported to bind to HER2 in cross-linking studies on human breast cancer cells. In addition, NDF has been shown to induce differentiation of human mammary tumor cells to milk-producing, growth-arrested cells, whereas the heregulin family have been reported to stimulate proliferation of cultured human breast cancers cell monolayers.

The means by which receptor polypeptides transduce regulatory signals in response to ligand binding is not fully understood, and continues to be the subject of intensive investigation. However, important components of the process have been uncovered, including the understanding that phosphorylation of and by cell surface receptors hold fundamental roles in signal transduction. In addition to the involvement of phosphorylation in the signal process, the intracellular phenomena of receptor dimerization and receptor crosstalk function as primary components of the circuit through which ligand binding triggers a resulting cellular response. Ligand binding to transmembrane receptor tyrosine kinases induces receptor dimerization, leading to activation of kinase function through the interaction of adjacent cytoplasmic domains. Receptor crosstalk refers to intracellular communication between two or more proximate receptor molecules mediated by, for example, activation of one receptor through a mechanism involving the kinase activity of the other. One particularly relevant example of such a phenomenon is the binding of EGF to the EGFR, resulting in activation of the EGFR kinase domain and cross-phosphorylation of HER2 (Kokai et al., 1989, Cell 58: 287-92; Stern et al., 1988, EMBO J. 7: 995-1001; King et al., 1989, Oncogene 4: 13-18).

### 3. SUMMARY OF THE INVENTION

HER4 is the fourth member of the EGFR-family of receptor tyrosine kinases and is likely to be involved not only in regulating normal cellular function but also in the loss of normal growth control associated with certain human cancers. In this connection, HER4 appears to be closely connected with certain carcinomas of epithelial origin, such as adenocarcinoma of the breast. As such, its discovery, and the elucidation of the HER4 coding sequence, open a number of novel approaches to the diagnosis and treatment of human cancers in which the aberrant expression and/or function of this cell surface receptor is involved.

The complete nucleotide sequence encoding the prototype HER4 polypeptide of the invention is disclosed herein, and provides the basis for several general aspects of the invention hereinafter described. Thus, the invention includes embodiments directly involving the production and use of HER4 polynucleotide molecules. In addition, the invention provides HER4 polypeptides, such as the prototype HER4 polypeptide disclosed and characterized in the sections which follow. Polypeptides sharing nearly equivalent structural characteristics with the prototype HER4 molecule are also included within the scope of this invention. Furthermore, the invention includes polypeptides which interact with HER4 expressed on the surface of certain cells thereby affecting their growth and/or differentiation. The invention is also directed to anti-HER4 antibodies, which have a variety of uses including but not limited to their use as components of novel

biological approaches to human cancer diagnosis and therapy provided by the invention.

The invention also relates to the discovery of an apparent functional relationship between HER4 and HER2, and the therapeutic aspects of the invention include those which are based on applicants' preliminary understanding of this relationship. Applicants' data strongly suggests that HER4 interacts with HER2 either by heterodimer formation or receptor cross-talk, and that such interaction appears to be one mechanism by which the HER4 receptor mediates effects on cell behavior. The reciprocal consequence is that HER2 activation is in some circumstances mediated through HER4.

#### 4. BRIEF DESCRIPTIONS OF THE FIGURES

FIG. 1. Nucleotide sequence [SEQ ID NO: 1] and deduced amino acid sequence [SEQ ID NO: 2] of HER4 (1308 amino acid residues). Nucleotides are numbered on the left, and amino acids are numbered above the sequence.

FIG. 2. Nucleotide sequence (FIG. 2(A) [SEQ ID NO: 3]; FIG. 2(B) [SEQ ID NO: 5] and deduced amino acid sequence (FIG. 2(A) [SEQ ID NO: 4]; FIG. 2(B) [SEQ ID NO: 6]) of cDNAs encoding HER4 variants. (A) HER4 with alternate 3' end and without autophosphorylation domain. This sequence is identical with that of HER4 shown in FIG. 1 up to nucleotide 3188, where the sequence diverges and the open reading frame stops after 13 amino acids, followed by an extended, unique 3'-untranslated region. (B) HER4 with N-terminal truncation. This sequence contains the 3'-portion of the HER4 sequence where nucleotide position 158 of the truncated sequence aligns with position 2335 of the complete HER4 sequence shown in FIG. 1 (just downstream from the region encoding the ATP-binding site of the HER4 kinase). The first 155 nucleotides of the truncated sequence are unique from HER4 and may represent the 5'-untranslated region of a transcript derived from a cryptic promoter within an intron of the HER4 gene. (Section 6.2.2., *infra*).

FIG. 3. The deduced amino acid sequence of two variant forms of human HER4 aligned with the full length HER4 receptor as represented in FIG. 1. Sequences are displayed using the single-letter code and are numbered on the right with the complete HER4 sequence on top and the variant sequences below. Identical residues are indicated by a colon between the aligned residues. (A) HER4 with alternate 3'-end, lacking an autophosphorylation domain [SEQ ID NO: 4]. This sequence is identical with that of HER4 [SEQ ID NO: 2] shown in FIG. 1 up to amino acid 1045, where the sequence diverges and continues for 13 amino acids before reaching a stop codon. (B) HER4 with N-terminal truncation [SEQ ID NO: 6]. This sequence is identical to the 3'-portion of the HER4 [SEQ ID NO: 2] shown in FIG. 1 beginning at amino acid 768. (Section 6.2.2., *infra*).

FIG. 4. Deduced amino acid sequence of human HER4 [SEQ ID NO: 2] and alignment with other human EGFR-family members (EGFR [SEQ ID NO: 7]; HER2 [SEQ ID NO: 8]; HER3 [SEQ ID NO: 9]). Sequences are displayed using the single-letter code and are numbered on the left. Identical residues are denoted with dots, gaps are introduced for optimal alignment, cysteine residues are marked with an asterisk, and N-linked glycosylation sites are denoted with a plus (+). Potential protein kinase C phosphorylation sites are indicated by arrows (HER4 amino acid positions 679, 685, and 699). The predicted ATP-binding site is shown with 4 circled crosses, C-terminal tyrosines are denoted with open triangles, and tyrosines in HER4 that are conserved with the major autophosphorylation sites in the EGFR are indicated with black triangles. The predicted extracellular domain extends from the boundary of the signal sequence marked by an arrow at position 25, to the hydrophobic transmembrane domain which is overlined from amino acid positions 650 through 675. Various subdomains are labeled on the right: I, II, III, and IV = extracellular subdomains (domains II and IV are cysteine-rich); TM = transmembrane domain; TK = tyrosine kinase domain. Domains I, III, TK are boxed.

FIG. 5. (A) Hydropathy profile of HER4, aligned with (B) Comparison of protein domains for HER4 (1308 amino acids), EGFR (1210 amino acids), HER2 (1255 amino acids), and HER3 (1342 amino acids). The signal peptide is represented by a stippled box, the cysteine-rich extracellular subdomains are hatched, the transmembrane domain is filled, and the cytoplasmic tyrosine kinase domain is stippled. The percent amino acid sequence identities between HER4 and other EGFR-family members are indicated. Sig, signal peptide; I, II, III, and IV, extracellular domains; TM, transmembrane domain; JM, juxtamembrane domain; Caln, calcium influx and internalization domain; 3'UTR, 3' untranslated region.

FIG. 6. Northern blot analysis of mRNA from human tissues hybridized to HER4 probes from (A) the 3'-autophosphorylation domain, and (B) the 5'-extracellular domain (see Section 6.2.3., *infra*). RNA size markers (in kilobases) are shown on the left. Lanes 1 through 8 represent 2 µg of poly(A)<sup>+</sup> mRNA from pancreas, kidney, skeletal muscle, liver, lung, placenta, brain, and heart, respectively.

FIG. 7. Immunoblot analysis of recombinant HER4 stably expressed in CHO-K1 cells, according to procedure outlined in Section 7.1.3, *infra*. Membrane preparations from CHO-K1 cells expressing recom-

binant HER4 were separated on 7% SDS-polyacrylamide gels and transferred to nitrocellulose. Blots were hybridized with (A) a monoclonal antibody to the C-terminus of HER2 (Ab3, Oncogene Science, Uniondale, NY) that cross-reacts with HER4 or (B) a sheep antipeptide polyclonal antibody to a common epitope of HER2 and HER4. Lane 1, parental CHO-K1 cells; lanes 2 - 4, CHO-K1/HER4 cell clones 6, 21, and 3, respectively. Note the 180 kDa HER4 protein and the 130 kDa cross-reactive species. The size in kilodaltons of prestained high molecular weight markers (BioRad, Richmond, CA) is shown on the left.

FIG. 8. Specific activation of HER4 tyrosine kinase by a breast cancer differentiation factor (see Section 8., *infra*). Four recombinant cell lines, each of which was engineered to overexpress a single member of EGFR-family of tyrosine kinase receptors (EGFR, HER2, HER3, and HER4), were prepared according to the methods described in Sections 7.1.2 and 8.1., *infra*. Cells from each of the four recombinant cell lines were stimulated with various ligand preparations and assayed for receptor tyrosine phosphorylation using the assay described in Section 8.2., *infra*. (A) CHO/HER4 #3 cells, (B) CHO/HER2 cells, (C) NRHER5 cells, and (D) 283/HER3 cells. Cells stimulated with: lane 1, buffer control; lane 2, 100 ng/ml EGF; lane 3, 200 ng/ml amphiregulin; lane 4, 10  $\mu$ l phenyl column fraction 17 (Section 9, *infra*); lane 5, 10  $\mu$ l phenyl column fraction 14 (Section 9., *infra*, and see description of FIG. 9 below). The size (in kilodaltons) of the prestained molecular weight markers are labeled on the left of each panel. The phosphorylated receptor in each series migrates just below the 221 kDa marker. Bands at the bottom of the gels are extraneous and are due to the reaction of secondary antibodies with the antibodies used in the immunoprecipitation.

FIG. 9. Biological and biochemical properties of the MDA-MB-453-cell differentiation activity purified from the conditioned media of HepG2 cells (Section 9., *infra*). (A, B, and C) Induction of morphologic differentiation. Conditioned media from HepG2 cells was subjected to ammonium sulfate fractionation, followed by dialysis against PBS. Dilutions of this material were added to MDA-MB-453 monolayer at the indicated protein concentrations. (A) control; (B) 80 ng per well; (C) 2.0  $\mu$ g per well. (D) Phenyl-5PW column elution profile monitored at 230 nm absorbance. (E) Stimulation of MDA-MB-453 tyrosine auto-phosphorylation with the following ligand preparations: None (control with no factor added); TGF- $\alpha$  (50 ng/ml); CM (16-fold concentrated HepG2 conditioned medium tested at 2  $\mu$ l and 10  $\mu$ l per well); fraction (phenyl column fractions 13 to 20, 10  $\mu$ l per well). (F) Densitometry analysis of the phosphorylation signals shown in (E).

FIG. 10. NDF-induced tyrosine phosphorylation of (A) MDA-MB-453 cells (lane 1, mock transfected COS cell supernatant; lane 2, NDF transfected COS cell supernatant); and (B) CHO/HER4 21-2 cells (lanes 1 and 2, mock transfected COS cell supernatant; lanes 3 and 4, NDF transfected COS cell supernatant). See Section 10., *infra*. Tyrosine phosphorylation was determined by the tyrosine kinase stimulation assay described in Section 8.2., *infra*.

FIG. 11. Regional location of the HER4 gene to human chromosome 2 band q33. (A) Distribution of 124 sites of hybridization on human chromosomes. (B) Distribution of autoradiographic grains on diagram of chromosome 2.

FIG. 12. Amino acid sequence of HER4-Ig fusion protein [SEQ ID NO: 10] (Section 5.4., *infra*).

## 5. DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed to HER4/p180<sup>erbB4</sup> ("HER4"), a closely related yet distinct member of the Human EGF Receptor (HER)/*neu* subfamily of receptor tyrosine kinases, as well as HER4-encoding polynucleotides (e.g., cDNAs, genomic DNAs, RNAs, anti-sense RNAs, etc.), the production of mature and precursor forms of HER4 from a HER4 polynucleotide coding sequence, recombinant HER4 expression vectors, HER4 analogues and derivatives, anti-HER4 antibodies, HER4 ligands, and diagnostic and therapeutic uses of HER4 polynucleotides, polypeptides, ligands, and antibodies in the field of human oncology and neurobiology.

The invention also reveals an apparent functional relationship between the HER4 and HER2 receptors involving HER4-mediated phosphorylation of HER2, potentially via intracellular receptor crosstalk or receptor dimerization. In this connection, the invention also provides a HER4 ligand capable of inducing cellular differentiation in breast carcinoma cells that appears to involve HER4-mediated phosphorylation of HER2. Furthermore, applicants' data provide evidence that NDF/HRG- $\alpha$  mediate biological effects on certain cells not solely through HER2, as has been reported in the literature, but instead by means of a direct interaction with HER4, or through an interaction with a HER2/HER4 complex. In cell lines expressing both HER2 and HER4, binding of NDF to HER4 may stimulate HER2 either by heterodimer formation of these two related receptors or by intracellular receptor crosstalk.

Unless otherwise indicated, the practice of the present invention utilizes standard techniques of molecular biology and molecular cloning, microbiology, immunology, and recombinant DNA known in the

art. Such techniques are described and explained throughout the literature, and can be found in a number of more comprehensive publications such as, for example, Maniatis et al, Molecular Cloning; A Laboratory Manual (Second Edition, 1989).

## 6 5.1. HER4 POLYNUCLEOTIDES

One aspect of the present invention is directed to HER4 polynucleotides, including recombinant polynucleotides encoding the prototype HER4 polypeptide shown in FIG. 1, polynucleotides which are related or are complementary thereto, and recombinant vectors and cell lines incorporating such recombinant polynucleotides. The term "recombinant polynucleotide" as used herein refers to a polynucleotide of genomic, cDNA, synthetic or semisynthetic origin which, by virtue of its origin or manipulation, is not associated with any portion of the polynucleotide with which it is associated in nature, and may be linked to a polynucleotide other than that to which it is linked in nature, and includes single or double stranded polymers of ribonucleotides, deoxyribonucleotides, nucleotide analogs, or combinations thereof. The term also includes various modifications known in the art, including but not limited to radioactive and chemical labels, methylation, caps, internucleotide modifications such as those with charged linkages (e.g., phosphorothioates, phosphorodithioates, etc.) and uncharged linkages (e.g., methyl phosphonates, phosphotriesters, phosphoamidites, carbamites, etc.), as well as those containing pendant moieties, intercalators, chelators, alkylators, etc. Related polynucleotides are those having a contiguous stretch of about 200 or more nucleotides and sharing at least about 80% homology to a corresponding sequence of nucleotides within the nucleotide sequence disclosed in FIG. 1. Several particular embodiments of such HER4 polynucleotides and vectors are provided in example Sections 6 and 7, *infra*.

HER4 polynucleotides may be obtained using a variety of general techniques known in the art, including molecular cloning and chemical synthetic methods. One method by which the molecular cloning of cDNAs encoding the prototype HER4 polypeptide of the invention (FIG. 1), as well as several HER4 polypeptide variants, is described by way of example in Section 6., *infra*. Conserved regions of the sequences of EGFR, HER2, HER3, and Xmrk are used for selection of the degenerate oligonucleotide primers which are then used to isolate HER4. Since many of these sequences have extended regions of amino acid identity, it is difficult to determine if a short PCR fragment represents a unique molecule or merely the species-specific counterpart of EGFR, HER2, or HER3. Often the species differences for one protein are as great as the differences within species for two distinct proteins. For example, fish Xmrk has regions of 47/55 (85%) amino acid identity to human EGFR, suggesting it might be the fish EGFR, however isolation of another clone that has an amino acid sequence identical to Xmrk in this region (57/57) shows a much higher homology to human EGFR in its flanking sequence (92% amino acid homology) thereby suggesting that it, and not Xmrk, is the fish EGFR (Wittbrodt, J. et al., 1989, Nature 342: 415-421). As described in Section 6., *infra*, it was necessary to confirm that a murine HER4/erbB4 PCR fragment was indeed a unique gene, and not the murine homolog of EGFR, HER2, or HER3, by isolating genomic fragments corresponding to murine EGFR, erbB2 and erbB3. Sequence analysis of these clones confirmed that this fragment was a novel member of the EGFR family. Notably a region of the murine clone had a stretch of 60/64 amino acid identity to human HER2, but comparison with the amino acid and DNA sequences of the other EGFR homologs from the same species (mouse) firmly established it encoded a novel transcript.

HER4 polynucleotides may be obtained from a variety of cell sources which produce HER4-like activities and/or which express HER4-encoding mRNA. In this connection, applicants have identified a number of suitable human cell sources for HER4 polynucleotides, including but not limited to brain, cerebellum, pituitary, heart, skeletal muscle, and a variety of breast carcinoma cell lines (see Section 6., *infra*).

For example, polynucleotides encoding HER4 polypeptides may be obtained by cDNA cloning from RNA isolated and purified from such cell sources or by genomic cloning. Either cDNA or genomic libraries of clones may be prepared using techniques well known in the art and may be screened for particular HER4-encoding DNAs with nucleotide probes which are substantially complementary to any portion of the HER4 gene. Various PCR cloning techniques may also be used to obtain the HER4 polynucleotides of the invention. A number of PCR cloning protocols suitable for the isolation of HER4 polynucleotides have been reported in the literature (see, for example, PCR protocols: A Guide to Methods and Applications, Eds. Inis et al., Academic Press, 1990).

For the construction of expression vectors, polynucleotides containing the entire coding region of the desired HER4 may be isolated as full length clones or prepared by splicing two or more polynucleotides together. Alternatively, HER4-encoding DNAs may be synthesized in whole or in part by chemical synthesis

using techniques standard in the art. Due to the inherent degeneracy of nucleotide coding sequences, any polynucleotide encoding the desired HER4 polypeptide may be used for recombinant expression. Thus, for example, the nucleotide sequence encoding the prototype HER4 of the invention provided in FIG. 1 may be altered by substituting nucleotides such that the same HER4 product is obtained.

5 The invention also provides a number of useful applications of the the HER4 polynucleotides of the invention, including but not limited to their use in the preparation of HER4 expression vectors, primers and probes to detect and/or clone HER4, and diagnostic reagents. Diagnostics based upon HER4 polynucleotides include various hybridization and PCR assays known in the art, utilizing HER4 polynucleotides as primers or probes, as appropriate. One particular aspect of the invention relates to a PCR kit comprising  
10 a pair of primers capable of priming cDNA synthesis in a PCR reaction, wherein each of the primers is a HER4 polynucleotide of the invention. Such a kit may be useful in the diagnosis of certain human cancers which are characterized by aberrant HER4 expression. For example, certain human carcinomas may overexpress HER4 relative to their normal cell counterparts, such as human carcinomas of the breast. Thus, detection of HER4 overexpression mRNA in breast tissue may be an indication of neoplasia. In another,  
15 related embodiment, human carcinomas characterized by overexpression of HER2 and expression or overexpression of HER4 may be diagnosed by a polynucleotide-based assay kit capable of detecting both HER2 and HER4 mRNAs, such a kit comprising, for example, a set of PCR primer pairs derived from divergent sequences in the HER2 and HER4 genes, respectively.

## 20 5.2. HER4 POLYPEPTIDES

Another aspect of the invention is directed to HER4 polypeptides, including the prototype HER4 polypeptide provided herein, as well as polypeptides derived from or having substantial homology to the amino acid sequence of the prototype HER4 molecule. The term "polypeptide" in this context refers to a  
25 polypeptide prepared by synthetic or recombinant means, or which is isolated from natural sources. The term "substantially homologous" in this context refers to polypeptides of about 80 or more amino acids sharing greater than about 90% amino acid homology to a corresponding contiguous amino acid sequence in the prototype HER4 primary structure (FIG. 1). The term "prototype HER4" refers to a polypeptide having the amino acid sequence of precursor or mature HER4 as provided in FIG. 1, which is encoded by  
30 the consensus cDNA nucleotide sequence also provided therein, or by any polynucleotide sequence which encodes the same amino acid sequence.

HER4 polypeptides of the invention may contain deletions, additions or substitutions of amino acid residues relative to the sequence of the prototype HER4 depicted in FIG. 1 which result in silent changes thus producing a bioactive product. Such amino acid substitutions may be made on the basis of similarity in  
35 polarity, charge, solubility, hydrophobicity, hydrophilicity and/or the amphipathic nature of the residues involved. For example, negatively charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine and arginine; amino acids with uncharged polar head groups or nonpolar head groups having similar hydrophilicity values include the following: leucine, isoleucine, valine; glycine, alanine; asparagine, glutamine; serine, threonine; phenylalanine, tyrosine.

40 The HER4 polypeptide depicted in FIG. 1 has all of the fundamental structural features characterizing the EGFR-family of receptor tyrosine kinases (Hanks et al., 1988, Science 241: 42-52). The precursor contains a single hydrophobic stretch of 26 amino acids characteristic of a transmembrane region that bisects the protein into a 625 amino acid extracellular ligand binding domain, and a 633 amino acid C-terminal cytoplasmic domain. The ligand binding domain can be further divided into 4 subdomains (I - IV),  
45 including two cysteine-rich regions (II, residues 186-334; and IV, residues 496-633), and two flanking domains (I, residues 29-185; and III, residues 335-495) that may define specificity for ligand binding (Lax et al., 1988, Mol. Cell. Biol. 8:1970-78). The extracellular domain of HER4 is most similar to HER3, where domains II-IV of HER4 share 58-87% identity to the respective domains of HER3. In contrast, the same regions of EGFR and HER2 exhibit 43-51% and 34-46% homology to HER4, respectively (FIG. 4). The 4  
50 extracellular subdomains of EGFR and HER2 share 39-50% identity. HER4 also conserves all 50 cysteines present in the extracellular portion of EGFR, HER2, and HER3, except that the HER2 protein lacks the fourth cysteine in domain IV. There are 11 potential N-linked glycosylation sites in HER4, conserving 4 of 12 potential sites in EGFR, 3 of 8 sites in HER2, and 4 of 10 sites in HER3.

Following the transmembrane domain of HER4 is a cytoplasmic juxtamembrane region of 37 amino  
55 acids. This region shares the highest degree of homology with EGFR (73% amino acid identity) and contains two consensus protein kinase C phosphorylation sites at amino acid residue numbers 679 (Serine) and 699 (Threonine) in the FIG. 1 sequence, the latter of which is present in EGFR and HER2. Notably, HER4 lacks a site analogous to Thr654 of EGFR. Phosphorylation of this residue in the EGFR appears to



block ligand-induced internalization and plays an important role in its transmembrane signaling (Livneh et al., 1988, Mol. Cell. Biol. 8: 2302-08). HER4 also contains Thr692 analogous to Thr694 of HER2. This threonine is absent in EGFR and HER3 and has been proposed to impart cell-type specificity to the mitogenic and transforming activity of the HER2 kinase (DiFiore et al. 1992, EMBO J. 11: 3927-33). The juxtamembrane region of HER4 also contains a MAP kinase consensus phosphorylation site at amino acid number 699 (Threonine), in a position homologous to Thr699 of EGFR which is phosphorylated by MAP kinase in response to EGF stimulation (Takishima et al., 1991, Proc. Natl. Acad. Sci. U.S.A. 88: 2520-25).

The remaining cytoplasmic portion of HER4 consists of a 276 amino acid tyrosine kinase domain, an acidic helical structure of 38 amino acids that is homologous to a domain required for ligand-induced internalization of the EGFR (Chen et al., 1989, Cell 59:33-43), and a 282 amino acid region containing 18 tyrosine residues characteristic of the autophosphorylation domains of other EGFR-related proteins (FIG. 4). The 276 amino acid tyrosine kinase domain conserves all the diagnostic structural motifs of a tyrosine kinase, and is most related to the catalytic domains of EGFR (78% identity) and HER2 (77% identity), and to a lesser degree, HER3 (63% identity). In this same region, EGFR and HER2 share 83% identity. Examples of the various conserved structural motifs include the following: the ATP-binding motif (GXGXXG) [SEQ ID NO: 11] with a distal lysine residue that is predicted to be involved in the phosphotransfer reaction (Hanks et al., 198, Science 241: 42-52; Hunter and Cooper, in The Enzymes Vol. 17 (eds. Boyer and Krebs) pp. 191-246 (Academic Press 1989)); tyrosine-kinase specific signature sequences (DLAARN [SEQ ID NO: 12] and PIKWMA [SEQ ID NO: 13]) and Tyr875 (FIG. 4), a residue that frequently serves as an autophosphorylation site in many tyrosine kinases (Hunter and Cooper, *supra*); and approximately 15 residues that are either highly or completely conserved among all known protein kinases (Flowman et al., 1990, Proc. Natl. Acad. Sci. U.S.A. 87: 4905-09; Hanks et al., *supra*). The C-terminal 282 amino acids of HER4 has limited homology with HER2 (27%) and EGFR (19%). However, the C-terminal domain of each EGFR-family receptor is proline-rich and conserves stretches of 2-7 amino acids that are generally centered around a tyrosine residue. These residues include the major tyrosine autophosphorylation sites of EGFR at Tyr1068, Tyr1086, Tyr1148, and Tyr1173 (FIG. 4, filled triangles; Margolis et al., 1989, J. Biol. Chem. 264: 10667-71).

### 5.3. RECOMBINANT SYNTHESIS OF HER4 POLYPEPTIDES

The HER4 polypeptides of the invention may be produced by the cloning and expression of DNA encoding the desired HER4 polypeptide. Such DNA may be ligated into a number of expression vectors well known in the art and suitable for use in a number of acceptable host organisms, in fused or mature form, and may contain a signal sequence to permit secretion. Both prokaryotic and eukaryotic host expression systems may be employed in the production of recombinant HER4 polypeptides. For example, the prototype HER4 precursor coding sequence or its functional equivalent may be used in a host cell capable of processing the precursor correctly. Alternatively, the coding sequence for mature HER4 may be used to directly express the mature HER4 molecule. Functional equivalents of the HER4 precursor coding sequence include any DNA sequence which, when expressed inside the appropriate host cell, is capable of directing the synthesis, processing and/or export of HER4.

Production of a HER4 polypeptide using recombinant DNA technology may be divided into a four-step process for the purposes of description: (1) isolation or generation of DNA encoding the desired HER4 polypeptide; (2) construction of an expression vector capable of directing the synthesis of the desired HER4 polypeptide; (3) transfection or transformation of appropriate host cells capable of replicating and expressing the HER4 coding sequence and/or processing the initial product to produce the desired HER4 polypeptide; and (4) identification and purification of the desired HER4 product.

#### 5.3.1. ISOLATION OR GENERATION OF HER4 ENCODING DNA

HER4-encoding DNA, or functional equivalents thereof, may be used to construct recombinant expression vectors which will direct the expression of the desired HER4 polypeptide product. In a specific embodiment, DNA encoding the prototype HER4 polypeptide (FIG. 1), or fragments or functional equivalents thereof, may be used to generate the recombinant molecules which will direct the expression of the recombinant HER4 product in appropriate host cells. HER4-encoding nucleotide sequences may be obtained from a variety of cell sources which produce HER4-like activities and/or which express HER4-encoding mRNA. For example, HER4-encoding cDNAs may be obtained from the breast adenocarcinoma cell line MDA-MB-453 (ATCC HTB131) as described in Section 6., *infra*. In addition, a number of human cell sources are suitable for obtaining HER4 cDNAs, including but not limited to various epidermoid and

breast carcinoma cells, and normal heart, kidney, and brain cells (see Section 6.2.3., *infra*).

The HER4 coding sequence may be obtained by molecular cloning from RNA isolated and purified from such cell sources or by genomic cloning. Either cDNA or genomic libraries of clones may be prepared using techniques well known in the art and may be screened for particular HER4-encoding DNAs with nucleotide probes which are substantially complementary to any portion of the HER4 gene. Alternatively, cDNA or genomic DNA may be used as templates for PCR cloning with suitable oligonucleotide primers. Full length clones, i.e., those containing the entire coding region of the desired HER4 may be selected for constructing expression vectors, or overlapping cDNAs can be ligated together to form a complete coding sequence. Alternatively, HER4-encoding DNAs may be synthesized in whole or in part by chemical synthesis using techniques standard in the art.

### 6.3.2. CONSTRUCTION OF HER4 EXPRESSION VECTORS

Various expression vector/host systems may be utilized equally well by those skilled in the art for the recombinant expression of HER4 polypeptides. Such systems include but are not limited to microorganisms such as bacteria transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing the desired HER4 coding sequence; yeast transformed with recombinant yeast expression vectors containing the desired HER4 coding sequence; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing the desired HER4 coding sequence; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing the desired HER4 coding sequence; or animal cell systems infected with recombinant virus expression vectors (e.g., adenovirus, vaccinia virus) including cell lines engineered to contain multiple copies of the HER4 DNA either stably amplified (e.g., CHO/dhfr, CHO/glutamine synthetase) or unstably amplified in double-minute chromosomes (e.g., murine cell lines).

The expression elements of these vectors vary in their strength and specificities. Depending on the host/vector system utilized, any one of a number of suitable transcription and translation elements may be used. For instance, when cloning in mammalian cell systems, promoters isolated from the genome of mammalian cells, (e.g., mouse metallothionein promoter) or from viruses that grow in these cells, (e.g., vaccinia virus 7.5K promoter or Moloney murine sarcoma virus long terminal repeat) may be used. Promoters produced by recombinant DNA or synthetic techniques may also be used to provide for transcription of the inserted sequences.

Specific initiation signals are also required for sufficient translation of inserted protein coding sequences. These signals include the ATG initiation codon and adjacent sequences. In cases where the entire HER4 gene including its own initiation codon and adjacent sequences are inserted into the appropriate expression vectors, no additional translational control signals may be needed. However, in cases where only a portion of the coding sequence is inserted, exogenous translational control signals, including the ATG initiation codon must be provided. Furthermore, the initiation codon must be in phase with the reading frame of the HER4 coding sequences to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of transcription attenuation sequences, enhancer elements, etc.

For example, in cases where an adenovirus is used as an expression vector, the desired HER4 coding sequence may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by *in vitro* or *in vivo* recombination. Insertion in a non-essential region of the viral genome (e.g., region E3 or E4) will result in a recombinant virus that is viable and capable of expressing HER4 in infected hosts. Similarly, the vaccinia 7.5K promoter may be used. An alternative expression system which could be used to express HER4 is an insect system. In one such system, Autographa californica nuclear polyhidrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. The HER4 coding sequence may be cloned into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter). Successful insertion of the HER4 coding sequence will result in inactivation of the polyhedrin gene and production of non-occluded recombinant virus (i.e., virus lacking the proteinaceous coat encoded by the polyhedrin gene). These recombinant viruses are then used to infect *Spodoptera frugiperda* cells in which the inserted gene is expressed. Yet another approach uses retroviral vectors prepared in amphotropic packaging cell lines, which permit high efficiency expression in numerous cell types. This method allows one to assess cell-type specific processing, regulation or function of the inserted protein coding sequence.

In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences; or modifies and processes the gene product in the specific fashion desired. Expression from certain promoters can be elevated in the presence of certain inducers. (e.g., zinc and cadmium ions for metallothionein promoters). Therefore, expression of the recombinant HER4 polypeptide may be controlled.

This is important if the protein product of the cloned foreign gene is lethal to host cells. Furthermore, modifications (e.g., phosphorylation) and processing (e.g., cleavage) of protein products are important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of protein. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed.

#### 5.3.3. TRANSFORMANTS EXPRESSING HER4 GENE PRODUCTS

The host cells which contain the recombinant coding sequence and which express the desired HER4 polypeptide product may be identified by at least four general approaches (a) DNA-DNA, DNA-RNA or RNA-antisense RNA hybridization; (b) the presence or absence of "marker" gene functions; (c) assessing the level of transcription as measured by the expression of HER4 mRNA transcripts in the host cell; and (d) detection of the HER4 product as measured by immunoassay and, ultimately, by its biological activities.

In the first approach, for example, the presence of HER4 coding sequences inserted into expression vectors can be detected by DNA-DNA hybridization using hybridization probes and/or primers for PCR reactions comprising polynucleotides that are homologous to the HER4 coding sequence.

In the second approach, the recombinant expression vector/host system can be identified and selected based upon the presence or absence of certain "marker" gene functions (e.g., thymidine kinase activity, resistance to antibiotics, resistance to methotrexate (MTX), resistance to methionine sulfoximine (MSX), transformation phenotype, occlusion body formation in baculovirus, etc.). For example, if the HER4 coding sequence is inserted within a marker gene sequence of the vector, recombinants containing that coding sequence can be identified by the absence of the marker gene function. Alternatively, a marker gene can be placed in tandem with the HER4 sequence under the control of the same or different promoter used to control the expression of the HER4 coding sequence. Expression of the marker in response to induction or selection indicates expression of the HER4 coding sequence. In a particular embodiment described by way of example herein, a HER4 expression vector incorporating glutamine synthetase as a selectable marker is constructed, used to transfect CHO cells, and amplified expression of HER4 in CHO cells is obtained by selection with increasing concentration of MSX.

In the third approach, transcriptional activity for the HER4 coding region can be assessed by hybridization assays. For example, polyadenylated RNA can be isolated and analyzed by Northern blot using a probe homologous to the HER4 coding sequence or particular portions thereof. Alternatively, total nucleic acids of the host cell may be extracted and assayed for hybridization to such probes.

In the fourth approach, the expression of HER4 can be assessed immunologically, for example by Western blots, immunoassays such as radioimmunoprecipitation, enzyme-linked immunoassays and the like. Alternatively, expression of HER4 may be assessed by detecting a biologically active product. Where the host cell secretes the gene product the cell free media obtained from the cultured transfectant host cell may be assayed for HER4 activity. Where the gene product is not secreted, cell lysates may be assayed for such activity. In either case, assays which measure ligand binding to HER4, HER4 phosphorylation, or other bioactivities of HER4 may be used.

#### 5.4. ANTI-HER4 ANTIBODIES

The invention is also directed to polyclonal and monoclonal antibodies which recognize epitopes of HER4 polypeptides. Anti-HER4 antibodies are expected to have a variety of useful applications in the field of oncology, several of which are described generally below. More detailed and specific descriptions of various uses for anti-HER4 antibodies are provided in the sections and subsections which follow. Briefly, anti-HER4 antibodies may be used for the detection and quantification of HER4 polypeptide expression in cultured cells, tissue samples, and *in vivo*. Such immunological detection of HER4 may be used, for example, to identify, monitor, and assist in the prognosis of neoplasms characterized by aberrant or attenuated HER4 expression and/or function. Additionally, monoclonal antibodies recognizing epitopes from different parts of the HER4 structure may be used to detect and/or distinguish between native HER4 and various subcomponent and/or mutant forms of the molecule. Anti-HER4 antibody preparations are also envisioned as useful biomodulatory agents capable of effectively treating particular human cancers. In addition to the various diagnostic and therapeutic utilities of anti-HER4 antibodies, a number of industrial

and research applications will be obvious to those skilled in the art, including, for example, the use of anti-HER4 antibodies as affinity reagents for the purification of HER4 polypeptides, and as immunological probes for elucidating the biosynthesis, metabolism and biological functions of HER4.

Anti-HER4 antibodies may be useful for influencing cell functions and behaviors which are directly or indirectly mediated by HER4. As an example, modulation of HER4 biological activity with anti-HER4 antibodies may influence HER2 activation and, as a consequence, modulate intracellular signals generated by HER2. In this regard, anti-HER4 antibodies may be useful to effectively block ligand-induced, HER4-mediated activation of HER2, thereby affecting HER2 biological activity. Conversely, anti-HER4 antibodies capable of acting as HER4 ligands may be used to trigger HER4 biological activity and/or initiate a ligand-induced, HER4-mediated effect on HER2 biological activity, resulting in a cellular response such as differentiation, growth inhibition, etc.

Additionally, anti-HER4 antibodies conjugated to cytotoxic compounds may be used to selectively target such compounds to tumor cells expressing HER4, resulting in tumor cell death and reduction or eradication of the tumor. In a particular embodiment, toxin-conjugated antibodies having the capacity to bind to HER4 and internalize into such cells are administered systemically for targeted cytotoxic effect. The preparation and use of radionuclide and toxin conjugated anti-HER4 antibodies are further described in Section 5.5., *infra*.

Overexpression of HER2 is associated with several human cancers. Applicants' data indicate that HER4 is expressed in certain human carcinomas in which HER2 overexpression is present. Therefore, anti-HER4 antibodies may have growth and differentiation regulatory effects on cells which overexpress HER2 in combination with HER4 expression, including but not limited to breast adenocarcinoma cells. Accordingly, this invention includes antibodies capable of binding to the HER4 receptor and modulating HER2 or HER2-HER4 functionality, thereby affecting a response in the target cell. For the treatment of cancers involving HER4-mediated regulation of HER2 biological activity, agents capable of selectively and specifically affecting the intracellular molecular interaction between these two receptors may be conjugated to internalizing anti-HER4 antibodies. The specificity of such agents may result in biological effects only in cells which co-express HER2 and HER4, such as breast cancer cells.

Various procedures known in the art may be used for the production of polyclonal antibodies to epitopes of HER4. For the production of polyclonal antibodies, a number of host animals are acceptable for the generation of anti-HER4 antibodies by immunization with one or more injections of a HER4 polypeptide preparation, including but not limited to rabbits, mice, rats, etc. Various adjuvants may be used to increase the immunological response in the host animal, depending on the host species, including but not limited to Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, oil emulsions, keyhole limpet hemocyanins, dinitrophenol, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and *Corynebacterium parvum*.

A monoclonal antibody to an epitope of HER4 may be prepared by using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include but are not limited to the hybridoma technique originally described by Kohler and Milstein (1975, *Nature* 256, 495-497), and the more recent human B-cell hybridoma technique (Kosbor et al., 1983, *Immunology Today* 4:72) and EBV-hybridoma technique (Cole et al., 1985, *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96). In addition techniques developed for the production of "chimeric antibodies" by splicing the genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity may be used (Morrison et al., 1984, *Proc. Natl. Acad. Sci.*, 81:6851-6855; Neuberger et al., 1984, *Nature*, 312:604-608; Takeda et al., 1985, *Nature*, 314:452-454). Alternatively, techniques described for the production of single chain antibodies (U.S. Patent 4,946,778) can be adapted to produce HER4-specific single chain antibodies. Recombinant human or humanized versions of anti-HER4 monoclonal antibodies are a preferred embodiment for human therapeutic applications. Humanized antibodies may be prepared according to procedures in the literature (e.g., Jones et al., 1986, *Nature* 321: 522-25; Reichman et al., 1988, *Nature* 332: 323-27; Verhoeven et al., 1988, *Science* 239: 1534-36). The recently described "gene conversion mutagenesis" strategy for the production of humanized anti-HER2 monoclonal antibody may also be employed in the production of humanized anti-HER4 antibodies (Carter et al., 1992, *Proc. Natl. Acad. Sci. U.S.A.* 89: 4285-89). Alternatively, techniques for generating a recombinant phage library of random combinations of heavy and light regions may be used to prepare recombinant anti-HER4 antibodies (e.g., Huse et al., 1989, *Science* 246: 1275-81).

As an example, anti-HER4 monoclonal antibodies may be generated by immunization of mice with cells selectively overexpressing HER4 (e.g., CHO/HER4 21-2 cells as deposited with the ATCC) or with partially purified recombinant HER4 polypeptides. In one embodiment, the full length HER4 polypeptide (FIG. 1)

may be expressed in Baculovirus systems, and membrane fractions of the recombinant cells used to immunize mice. Hybridomas are then screened on CHO/HER4 cells (e.g., CHO HER4 21-2 cells as deposited with the ATCC) to identify monoclonal antibodies reactive with the extracellular domain of HER4. Such monoclonal antibodies may be evaluated for their ability to block NDF, or HepG2-differentiating factor, binding to HER4; for their ability to bind and stay resident on the cell surface, or to internalize into cells expressing HER4; and for their ability to directly upregulate or downregulate HER4 tyrosine autophosphorylation and/or to directly induce a HER4-mediated signal resulting in modulation of cell growth or differentiation. In this connection, monoclonal antibodies N28 and N29, directed to HER2, specifically bind HER2 with high affinity. However, monoclonal N29 binding results in receptor internalization and down-regulation, morphologic differentiation, and inhibition of HER2 expressing tumor cells in athymic mice. In contrast, monoclonal N28 binding to HER2 expressing cells results in stimulation of autophosphorylation, and an acceleration of tumor cell growth both *in vitro* and *in vivo* (Bacus et al., 1992, Cancer Res. 52: 2580-89; Stancovski et al., 1991, Proc. Natl. Acad. Sci. U.S.A. 88: 8691-95). In yet another embodiment, a soluble recombinant HER4-Immunoglobulin (HER4-Ig) fusion protein is expressed and purified on a Protein A affinity column. The amino acid sequence of one such HER4-Ig fusion protein is provided in FIG. 12. The soluble HER4-Ig fusion protein may then be used to screen phage libraries designed so that all available combinations of a variable domain of the antibody binding site are presented on the surfaces of the phages in the library. Recombinant anti-HER4 antibodies may be propagated from phage which specifically recognize the HER4-Ig fusion protein.

Antibody fragments which contain the idiotype of the molecule may be generated by known techniques. For example, such fragments include but are not limited to: the F(ab')<sub>2</sub> fragment which can be produced by pepsin digestion of the intact antibody molecule; the Fab' fragments which can be generated by reducing the disulfide bridges of the F(ab')<sub>2</sub> fragment, and the two Fab fragments which can be generated by treating the antibody molecule with papain and a reducing agent. Alternatively, Fab expression libraries may be constructed (Huse et al., 1989, Science, 246:1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity to HER4 protein.

## 5.5. DIAGNOSTIC METHODS

The invention also relates to the detection of human neoplastic conditions, particularly carcinomas of epithelial origin, and more particularly human breast carcinomas. In one embodiment, oligomers corresponding to portions of the consensus HER4 cDNA sequence provided in FIG. 1 are used for the quantitative detection of HER4 mRNA levels in a human biological sample, such as blood, serum, or tissue biopsy samples, using a suitable hybridization or PCR format assay, in order to detect cells or tissues expressing abnormally high levels of HER4 as an indication of neoplasia. In a related embodiment, detection of HER4 mRNA may be combined with the detection HER2 mRNA overexpression, using appropriate HER2 sequences, to identify neoplasias in which a functional relationship between HER2 and HER4 may exist.

In another embodiment, labeled anti-HER4 antibodies or antibody derivatives are used to detect the presence of HER4 in biological samples, using a variety of immunoassay formats well known in the art, and may be used for *in situ* diagnostic radioimmunoimaging. Current diagnostic and staging techniques do not routinely provide a comprehensive scan of the body for metastatic tumors. Accordingly, anti-HER4 antibodies labeled with, for example, fluorescent, chemiluminescent, and radioactive molecules may overcome this limitation. In a preferred embodiment, a gamma-emitting diagnostic radionuclide is attached to a monoclonal antibody which is specific for an epitope of HER4, but not significantly cross-reactive with other EGFR-family members. The labeled antibody is then injected into a patient systemically, and total body imaging for the distribution and density of HER4 molecules is performed using gamma cameras, followed by localized imaging using computerized tomography or magnetic resonance imaging to confirm and/or evaluate the condition, if necessary. Preferred diagnostic radionuclides include but are not limited to technetium-99m, indium-111, iodine-123, and iodine-131.

Recombinant antibody-metallothionein chimeras (Ab-MTs) may be generated as recently described (Das et al., 1992, Proc. Natl. Acad. Sci. U.S.A. 89: 9749-53). Such Ab-MTs can be loaded with technetium-99m by virtue of the metallothionein chelating function, and may offer advantages over chemically conjugated chelators. In particular, the highly conserved metallothionein structure may result in minimal immunogenicity.

## 5.6. TARGETED CANCER THERAPY

The invention is also directed to methods for the treatment of human cancers involving abnormal expression and/or function of HER4 and cancers in which HER2 overexpression is combined with the proximate expression of HER4, including but not limited to human breast carcinomas and other neoplasms overexpressing HER4 or overexpressing HER2 in combination with expression of HER4. The cancer therapy methods of the invention are generally based on treatments with unconjugated, toxin- or radionuclide-conjugated HER4 antibodies, ligands, and derivatives or fragments thereof. In one specific embodiment, such HER4 antibodies may be used for systemic and targeted therapy of certain cancers overexpressing HER2 and/or HER4, such as metastatic breast cancer, with minimal toxicity to normal tissues and organs. Importantly, in this connection, an anti-HER2 monoclonal antibody has been shown to inhibit the growth of human tumor cells overexpressing HER2 (Bacus et al., 1992, Cancer Res. 52: 2580-89). In addition to conjugated antibody therapy, modulation of NDF signaling through HER4 may provide a means to affect the growth and differentiation of cells overexpressing HER2, such as certain breast cancer cells, using HER4-neutralizing monoclonal antibodies, NDF/HER4 antagonists, monoclonal antibodies or ligands which act as super-agonists for HER4 activation, or agents which block the interaction between HER2 and HER4, either by disrupting heterodimer formation or by blocking HER-mediated phosphorylation of the HER2 substrate.

For targeted immunotoxin-mediated cancer therapy, various drugs or toxins may be conjugated to anti-HER4 antibodies and fragments thereof, such as plant and bacterial toxins. For example, ricin, a cytotoxin from the *Ricinus communis* plant may be conjugated to an anti-HER4 antibody using methods known in the art (e.g., Blakey et al., 1988, Prog. Allergy 45: 50-90; Marsh and Neville, 1988, J. Immunol. 140: 3674-78). Once ricin is inside the cell cytoplasm, its A chain inhibits protein synthesis by inactivating the 60S ribosomal subunit (May et al., 1989, EMBO J. 8: 301-08). Immunotoxins of ricin are therefore extremely cytotoxic. However, ricin immunotoxins are not ideally specific because the B chain can bind to virtually all cell surface receptors, and immunotoxins made with ricin A chain alone have increased specificity. Recombinant or deglycosylated forms of the ricin A chain may result in improved survival (i.e., slower clearance from circulation) of the immunotoxins. Methods for conjugating ricin A chain to antibodies are known (e.g., Vitella and Thorpe, in: Seminars in Cell Biology, pp47-58; Saunders, Philadelphia 1991). Additional toxins which may be used in the formulation of immunotoxins include but are not limited to daunorubicin, methotrexate, ribosome inhibitors (e.g., trichosanthin, trichokirin, gelonin, saporin, mormordin, and pokeweed antiviral protein) and various bacterial toxins (e.g., *Pseudomonas* endotoxin). Immunotoxins for targeted cancer therapy may be administered by any route which will result in antibody interaction with the target cancer cells, including systemic administration and injection directly to the site of tumor.

For targeted radiotherapy using anti-HER4 antibodies, preferred radionuclides for labeling include alpha, beta, and Auger electron emitters. Examples of alpha emitters include astatine 211 and bismuth 212; beta emitters include iodine 131, rhenium 188, copper 67 and yttrium 90; and iodine 125 is an example of an Auger electron emitter.

## 5.7. ASSAYS FOR THE IDENTIFICATION OF HER4 LIGANDS

Cell lines overexpressing a single member of the EGFR-family can be generated by transfection of a variety of parental cell types with an appropriate expression vector as described in section 7., *infra*. Candidate ligands, or partially purified preparations, may be applied to such cells and assayed for receptor binding and/or activation. For example, a CHO-K1 cell line transfected with a HER4 expression plasmid and lacking detectable EGFR, HER2, or HER3 may be used to screen for HER4-specific ligands. A particular embodiment of such a cell line is described in Section 7., *infra* and has been deposited with the ATCC (CHO/HER4 21-2). Ligands may be identified by detection of HER4 autophosphorylation, stimulation of DNA synthesis, induction of morphologic differentiation, relief from serum or growth factor requirements in the culture media, and direct binding of labeled purified growth factor. The invention also relates to a bioassay for testing potential analogs of HER4 ligands based on a capacity to affect a biological activity mediated by the HER4 receptor.

## 5.8 HER4 ANALOGUES

The production and use of derivatives, analogues and peptides related to HER4 are also envisioned and are within the scope of the invention. Such derivatives, analogues and peptides may be used to compete with native HER4 for binding of HER4 specific ligand, thereby inhibiting HER4 signal transduction and function. The inhibition of HER4 function may be utilized in several applications, including but not limited to

the treatment of cancers in which HER4 biological activity is involved.

In a specific embodiment, a series of deletion mutants in the HER4 nucleotide coding sequence depicted in FIG.1 may be constructed and analyzed to determine the minimum amino acid sequence requirements for binding of a HER4 ligand. Deletion mutants of the HER4 coding sequence may be constructed using methods known in the art which include but are not limited to use of nucleases and/or restriction enzymes; site-directed mutagenesis techniques, PCR, etc. The mutated polypeptides expressed may be assayed for their ability to bind HER4 ligand.

The DNA sequence encoding the desired HER4 analogue may then be cloned into an appropriate expression vector for overexpression in either bacteria or eukaryotic cells. Peptides may be purified from cell extracts in a number of ways including but not limited to ion-exchange chromatography or affinity chromatography using HER4 ligand or antibody. Alternatively, polypeptides may be synthesized by solid phase techniques followed by cleavage from resin and purification by high performance liquid chromatography.

## 6. EXAMPLE: ISOLATION OF cDNAs ENCODING HER4

EGFR and the related proteins, HER2, HER3, and Xmrk exhibit extensive amino acid homology in their tyrosine kinase domains (Kaplan et al., 1991, Nature 350: 158-160; Wen et al., 1992, Cell 69: 559-72; Holmes et al., 1992, Science 256: 1205-10; Hirai et al., 1987, Science 238: 1717-20). In addition, there is strict conservation of the exon-intron boundaries within the genomic regions that encode these catalytic domains (Wen et al., *supra*; Lindberg and Hunter, 1990, Mol. Cell. Biol. 10: 6316-24; and unpublished observations). Degenerate oligonucleotide primers were designed based on conserved amino acids encoded by a single exon or adjacent exons from the kinase domains of these four proteins. These primers were used in a polymerase chain reaction (PCR) to isolate genomic fragments corresponding to murine EGFR, erbB2 and erbB3. In addition, a highly related DNA fragment (designated MER4) was identified as distinct from these other genes. A similar strategy was used to obtain a cDNA clone corresponding to the human homologue of MER4 from the breast cancer cell line, MDA-MB-453. Using this fragment as a probe, several breast cancer cell lines and human heart were found to be an abundant source of the EGFR-related transcript. cDNA libraries were constructed using RNA from human heart and MDA-MB-453 cells, and overlapping clones were isolated spanning the complete open reading frame of HER4/erbB4.

### 6.1. MATERIALS AND METHODS

#### 6.1.1. MOLECULAR CLONING

Several pools of degenerate oligonucleotides were synthesized based on conserved sequences from EGFR-family members (Table I).

5'-ACNGTNTGGGARYTNAYHAC-3' [SEQ ID NO: 14]; 5'-CAYGTNAARATHACNGAYTTYGG-3' [SEQ ID NO: 15]; 5'-GACGAATTCNATHAARTGGATGGC [SEQ ID NO: 16]; 5'-ACAYTTNARDATDATCATRTANAC-3' [SEQ ID NO: 17]; 5'-AANGTCATNARYTCCCA-3' [SEQ ID NO: 18]; 5'-TCCAGNGCGATCCAYTT-DATNGG-3' [SEQ ID NO: 19]; 5'-GGRTCDATCATCCARCCT-3' [SEQ ID NO: 20]; 5'-CTGCTGTGAGCATC-GATCAT-3' [SEQ ID NO: 21]; TWELMT [SEQ ID NO: 22]; HVKITDFG [SEQ ID NO: 23]; PIKWMA [SEQ ID NO: 13]; VYMIILK [SEQ ID NO: 24]; WELMTF [SEQ ID NO: 25]; PIKWMALE [SEQ ID NO: 26]; CWMIDP [SEQ ID NO: 27]

Total genomic DNA was isolated from subconfluent murine K1735 melanoma cells and used as a template with these oligonucleotide primers in a 40 cycle PCR amplification. PCR products were resolved on agarose gels and hybridized to <sup>32</sup>P-labeled probes from the kinase domain of human EGFR and HER2. Distinct DNA bands were isolated and subcloned for sequence analysis. Using the degenerate oligonucleotides H4VWELM and H4VYMIL as primers in a PCR amplification (Plowman et al., 1990, Proc. Natl. Acad. Sci. U.S.A. 87: 4905-09), one clone (MER4-85) was identified that contained a 144 nucleotide insert corresponding to murine erbB4. This <sup>32</sup>P-labeled insert was used to isolate a 17-kilobase fragment from a murine T-cell genomic library (Stratagene, La Jolla, CA) that was found to contain two exons of the murine erbB4 gene. A specific oligonucleotide (4M3070) was synthesized based on the DNA sequence of an erbB4 exon, and used in a PCR protocol with a degenerate 5'-oligonucleotide (H4PIKWMA) on a template of single stranded MDA-MB-453 cDNA. This reaction generated a 260 nucleotide fragment (pMDAPIK) corresponding to human HER4. cDNA libraries were constructed in lambda ZAP II (Stratagene) from oligo(dT)- and specific-primed MDA-MB453 and human heart RNA (Plowman et al., *supra*; Plowman et al., 1990, Mol. Cell. Biol. 10: 1969-81). HER4-specific clones were isolated by probing the libraries with the

<sup>32</sup>P-labeled insert from pMDAPIK. To complete the cloning of the 5'-portion of HER4, we used a PCR strategy to allow for rapid amplification of cDNA ends (Plowman et al., *supra*; Frohman et al., 1988, Proc. Natl. Acad. Sci. U.S.A. 85: 8988-8992). All cDNA clones and several PCR generated clones were sequenced on both strands using T7 polymerase with oligonucleotide primers (Tabor and Richardson, 1987, Proc. Natl. Acad. Sci. U.S.A. 84: 4767-71).

**TABLE I**  
**OLIGONUCLEOTIDE PREPARATIONS FOR CLONING HER4**

Designation	Nucleotide Sequence <sup>1</sup>	Degeneracy	Encoded Sequence	Orientation
H4TVWELM	5'-ACNGTNTGGGARYTNAYHAC-3'	256-fold	TVWELMT	sense
H4KITDFG	5'-CAYGTNAARATHACNGAYTTYGG-3'	768-fold	HVKITDFG	sense
H4PIKMA	5'-GACGAATTCNATHAARTGGATGGC	48-fold	PIKMA	sense
H4VYMIIL	5'-ACAYTTNARDATDATCATRTANAC-3'	576-fold	VYMIILK	antisense
H4WELMTF	5'-AANGTCATNARYTCCCA-3'	32-fold	WELMTF	antisense
H4PIKMA	5'-TCCAGNCCGATCCAYTTDATNGG-3'	96-fold	PIKMALE	antisense
H4CWMIDP	5'-GGRTCDATCATCCARCT-3'	12-fold	CWMIDP	antisense
4H3070	5'-CTGCTGTCAGCATCGATCAT-3'	zero	erbB4 exon	antisense

<sup>1</sup>Degenerate nucleotide residue designations:

D = A, G, or T;  
H = A, C, or T;  
N = A, C, G, or T;  
R = A or G; and  
Y = C or T.

#### 6.1.2. NORTHERN BLOT ANALYSIS

3'- and 5'-HER4 specific [ $\alpha$ -<sup>32</sup>P]UTP-labeled antisense RNA probes were synthesized from the linearized plasmids pH1B1.6 (containing an 800 bp HER4 fragment beginning at nucleotide 3098) and p5'H4E7 (containing a 1 kb fragment from the 5'-end of the HER4 sequence), respectively. For tissue distribution analysis (Section 6.2.2., *infra*), the Northern blot (Clontech, Palo Alto, CA) contained 2  $\mu$ g poly(A)<sup>+</sup> mRNA per lane from 8 human tissue samples immobilized on a nylon membrane. The filter was prehybridized at 60°C for several hours in RNA hybridization mixture (50% formamide, 5XSSC, 0.5% SDS, 10X Denhardt's solution, 100  $\mu$ g/ml denatured herring sperm DNA, 100  $\mu$ g/ml tRNA, and 10  $\mu$ g/ml polyadenosine) and hybridized in the same buffer at 60°C, overnight with 1-1.5 x 10<sup>6</sup> cpm/ml of <sup>32</sup>P-labeled antisense RNA probe. The filters were washed in 0.1XSSC/0.1% SDS, 65°C, and exposed overnight on a phosphorimager (Molecular Dynamics, Sunnyvale, CA).

#### 6.1.3. SEMI-QUANTITATIVE PCR DETECTION OF HER4

RNA was isolated from a variety of human cell lines, fresh frozen tissues, and primary tumors. Single stranded cDNA was synthesized from 10  $\mu$ g of each RNA by priming with an oligonucleotide containing a T<sub>17</sub> track on its 3'-end (XSCT17:5'-GACTCGAGTCGACATCGATTTTTTTTTTTTTTTT-3') (SEQ ID NO: 28). 1% or 5% of each single strand template preparation was then used in a 35 cycle PCR reaction with two HER4-specific oligonucleotides: 4H2874: 5'-GAAGAAAGACGACTCGTTCATCGG-3', [SEQ ID NO: 29], and 4H2965: 5'-GACCATGACCATGTAAACGTCAATA-3') [SEQ ID NO: 30]. Reaction products were electrophoresed on 2% agarose gels, stained with ethidium bromide and photographed on a UV light box. The relative intensity of the 291-bp HER4-specific bands were estimated for each sample as shown in Table II.



### 6.2.1. SEQUENCE ANALYSIS OF cDNA CLONES ENCODING HER4

cDNA clones encoding parts of the HER4 coding and non-coding nucleotide sequences were isolated by PCR cloning according to the method outlined in Section 6.1.1., *supra*. The complete HER4 nucleotide sequence assembled from these cDNAs is shown in FIG. 1 and contains a single open reading frame encoding a polypeptide of 1308 amino acids. The HER4 coding region is flanked by a 33 nucleotide 5'-untranslated region and a 1517 nucleotide 3'-untranslated region ending with a poly(A) tail. A 25 amino acid hydrophobic signal sequence follows a consensus initiating methionine at position number 1 in the amino acid sequence depicted in FIG.1. In relation to this signal sequence, the mature HER4 polypeptide would be predicted to begin at amino acid residue number 28 in the sequence depicted in FIG. 1 (Gln), followed by the next 1283 amino acids in the sequence. Thus the prototype mature HER4 of the invention is a polypeptide of 1284 amino acids, having a calculated Mr of 144,280 daltons and an amino acid sequence corresponding to residues 28 through 1308 in FIG. 1.

Comparison of the HER4 nucleotide and deduced amino acid sequences (FIG. 1) with the available DNA and protein sequence databases indicated that the HER4 nucleotide sequence is unique, and revealed a 60/64 amino acid identity with HER2 and a 54/54 amino acid identity to a fragment of a rat EGFR homolog, tyro-2.

### 6.2.2. SEQUENCE ANALYSIS OF RELATED cDNAs

Several cDNAs encoding polypeptides related to the prototype HER4 polypeptide (FIG. 1) were also isolated from the MDA-MB-453 cDNA library and comprised two forms.

The first alternative type of cDNA was identical to the consensus HER4 nucleotide sequence up to nucleotide 3168 (encoding Arg at amino acid position 1045 in the FIG. 1 sequence) and then abruptly diverges into an apparently unrelated sequence (FIG. 2A, FIG. 3A). Downstream from this residue the open reading frame continues for another 13 amino acids before reaching a stop codon followed by a 2 kb 3'-untranslated sequence and poly(A) tail. This cDNA would be predicted to result in a HER4 variant having the C-terminal autophosphorylation domain of the prototype HER4 deleted.

A second type of cDNA was isolated as 4 independent clones each with a 3'-sequence identical to the HER4 consensus, but then diverging on the 5'-side of nucleotide 2335 (encoding Glu at amino acid position 768 in the FIG. 1 sequence), continuing upstream for only another 114-154 nucleotides (FIG. 2B, FIG. 3B). Nucleotide 2335 is the precise location of an intron-exon junction in the HER2 gene (Coussens et al., 1985, *Science* 230: 1132-39; Semba et al., 1985, *Proc. Natl. Acad. Sci. U.S.A.* 82: 6497-6501), suggesting these cDNAs could be derived from mRNAs that have initiated from a cryptic promoter within the flanking intron. These 5'-truncated transcripts contain an open reading frame identical to that of the HER4 cDNA sequence of FIG. 1, beginning with the codon for Met at amino acid position 772 in FIG. 1. These cDNAs would be predicted to encode a cytoplasmic HER4 variant polypeptide that initiates just downstream from the ATP-binding domain of the HER4 kinase.

### 6.2.3. HUMAN TISSUE DISTRIBUTION OF HER4 EXPRESSION

Northern blots of poly(A)<sup>+</sup> mRNA from human tissue samples were hybridized with antisense RNA probes to the 3'-end of HER4, encoding the autophosphorylation domain, as described in Section 6.1.2., *supra*. A HER4 mRNA transcript of approximately 6kb was identified, and was found to be most abundant in the heart and skeletal muscle (FIG. 6A). An mRNA of greater than approximately 15 kb was detected in the brain, with lower levels also detected in heart, skeletal muscle, kidney, and pancreas tissue samples.

The same blot was stripped and rehybridized with a probe from the 5'-end of HER4, within the extracellular domain coding region, using identical procedures. This hybridization confirmed the distribution of the 15 kb HER4 mRNA species, and detected a 6.5 kb mRNA species in heart, skeletal muscle, kidney, and pancreas tissue samples (FIG. 6B) with weaker signals in lung, liver, and placenta. In addition, minor transcripts of 1.7-2.8 kb were also detected in pancreas, lung, brain, and skeletal muscle tissue samples. The significance of the different sized RNA transcripts is not known.

Various human tissues were also examined for the presence of HER4 mRNA using the semi-quantitative PCR assay described in Section 6.1.3., *supra*. The results are shown in Table II, together with results of the assay on primary tumor samples and neoplastic cell lines (Section 6.2.4., immediately below). These results correlate well with the Northern and solution hybridization analysis results on the selected RNA samples. The highest levels of HER4 transcript expression were found in heart, kidney, and brain tissue samples. In addition, high levels of HER4 mRNA expression were found in parathyroid, cerebellum,

pituitary, spleen, testis, and breast tissue samples. Lower expression levels were found in thymus, lung, salivary gland, and pancreas tissue samples. Finally, low or negative expression was observed in liver, prostate, ovary, adrenal, colon, duodenum, epidermis, and bone marrow samples.

#### 5 **6.2.4. HER4 mRNA EXPRESSION IN PRIMARY TUMORS AND VARIOUS CELL LINES OF NEOPLASTIC ORIGIN**

HER4 mRNA expression profiles in several primary tumors and a number of cell lines of diverse neoplastic origin were determined with the semi-quantitative PCR assay (Section 6.1.3, *supra*) using primers from sequences in the HER4 kinase domain. The results are included in Table II. This analysis detected the highest expression of HER4 RNA in 4 human mammary adenocarcinoma cell lines (T-47D, MDA-MB-453, BT-474, and H3398), and in neuroblastoma (SK-N-MC), and pancreatic carcinoma (Hs768T) cell lines. Intermediate expression was detected in 3 additional mammary carcinoma cell lines (MCF-7, MDA-MB-330, MDA-MB-361). Low or undetectable expression was found in other cell lines derived from carcinomas of the breast (MDA-MB-231, MDA-MB-157, MDA-MB-468, SK-BR-3), kidney (Caki-1, Caki-2, G-401), liver (SK-HEP-1, HepG2), pancreas (PANC-1, AsPC-1, Capan-1), colon (HT-29), cervix (CaSki), vulva (A-41), ovary (PA-1, Caov-3), melanoma (SK-MEL-28), or in a variety of leukemic cell lines. Finally, high level expression was observed in Wilms (kidney) and breast carcinoma primary tumor samples.

**TABLE II**  
**HER4 EXPRESSION BY PRC ANALYSIS**

<u>VERY STRONG</u>	<u>STRONG</u>	<u>MEDIUM</u>
T47D (br ast)	MDA-MB-453 (breast) BT-474 (breast) H3396 (breast) Hs766T (pancreatic) SK-N-MC (neural) Wilms Tumor(kidney)	MCF-7 (breast) MDA-MB-330 (breast) MDA-MB-157 (breast) JEG-3 (choriocarcinoma) HEPM (palate) 458 (medullablastoma) Breast Carcinoma
Kidney Heart Parathyroid	Brain Cerebellum Pituitary Breast Testis Spleen	Skeletal Muscle Thymus Pancreas Lung Salivary Gland
<u>WEAK</u>	<u>NEGATIVE</u>	
MDA-MB-231 (breast) MDA-MB-157 (breast) SK-BR-3 (breast) A-431 (vulva) Caki-1 (kidney) Caki-2 (kidney) SK-HEP-1 (liver) THP-1 (macrophage)	MDA-MB-468 (breast) G-401 (kidney) HepG2 (liver) PANC-1 (pancreas) AsPC-1 (pancreas) Capan-1 (pancreas) HT-29 (colon) CaSki (cervix) PA-1 (ovary) Caov-3 (ovary) SK-MEL-28 (melanoma) HUF (fibroblast) H2981 (lung) Ovarian tumor GEO (colon) ALL bone marrow AML bone marrow Duodenum Epidermis Liver Bone marrow stroma	
Prostate Adrenal Ovary Colon Placenta		

## 7. EXAMPLE: RECOMBINANT EXPRESSION OF HER4

### 7.1. MATERIALS AND METHODS

#### 7.1.1. CHO-KI CELLS AND CULTURE CONDITIONS

CHO-KI cells were obtained from the ATCC (Accession Number CCL 61). These cells lack any detectable EGFR, HER2, or HER3 by immunoblot, tyrosine phosphorylation, and <sup>35</sup>S-labeled immunoprecipitation analysis. Transfected cell colonies expressing HER4 were selected in glutamine-free Glasgow modified Eagle's medium (GMEM-S, Gibco) supplemented with 10% dialyzed fetal bovine serum an increasing concentrations of methionine sulfoximine (Bebbington, 1991, in Methods: A Companion to Methods in Enzymology 2: 136-145 Academic Press).

## 7.1.2. EXPRESSION VECTOR CONSTRUCTION AND TRANSFECTIONS

The complete 4 kilobase coding sequence of prototype HER4 was reconstructed and inserted into a glutamine synthetase expression vector, pEE14, under the control of the cytomegalovirus immediate-early promoter (Bebbington, *supra*) to generate the HER4 expression vector pEEHER4. This construct (pEEHER4) was linearized with *Mlu*I and transfected into CHO-K1 cells by calcium phosphate precipitation using standard techniques. Cells were placed on selective media consisting of GMEW-S supplemented with 10% dialyzed fetal bovine serum and methionine sulfoximine at an initial concentration of 25  $\mu$ M (L-MSX) as described in Bebbington, *supra*, for the selection of initial resistant colonies. After 2 weeks, isolated colonies were transferred to 48-well plates and expanded for HER4 expression immunoassays as described immediately below. Subsequent rounds of selection using higher concentrations of MSX were used to isolate cell colonies tolerating the highest concentrations of MSX. A number of CHO/HER4 clones selected at various concentrations of MSX were isolated in this manner.

### 7.1.3. HER4 EXPRESSION IMMUNOASSAY

Confluent cell monolayers were scraped into hypotonic lysis buffer (10 mM Tris pH7.4, 1 mM KCl, 2 mM MgCl<sub>2</sub>) at 4°C, dounce homogenized with 30 strokes, and the cell debris was removed by centrifugation at 3500 x g, 5 min. Membrane fractions were collected by centrifugation at 100,000 x g, 20 min, and the pellet was resuspended in hot Laemmli sample buffer with 2-mercaptoethanol. Expression of the HER4 polypeptide was detected by immunoblot analysis on solubilized cells or membrane preparations using HER2 immunoreagents generated to either a 19 amino acid region of the HER2 kinase domain, which coincidentally is identical to the HER4 sequence (residues 927-945), or to the C-terminal 14 residues of HER2, which share a stretch of 7 consecutive residues with a region near the C-terminus of HER4. On further amplification, HER4 was detected from solubilized cell extracts by immunoblot analysis with PY20 anti-phosphotyrosine antibody (ICN Biochemicals), presumably reflecting autoactivation and auto-phosphorylation of HER4 due to receptor aggregation resulting from abberantly high receptor density. More specifically, expression was detected by immunoblotting with a primary murine monoclonal antibody to HER2 (Neu-Ab3, Oncogene Science) diluted 1:50 in blotto (2.5% dry milk, 0.2% NP40 in PBS) using <sup>125</sup>I-goat anti-mouse Ig F(ab')<sub>2</sub> (Amersham, UK) diluted 1:500 in blotto as a second antibody. Alternatively, a sheep polyclonal antipeptide antibody against HER2 residues 929-947 (Cambridge Research Biochemicals, Valleystream, NY) was used as a primary immunoreagent diluted 1:100 in blotto with <sup>125</sup>I-Protein G (Amersham) diluted 1:200 in blotto as a second antibody. Filters were washed with blotto and exposed overnight on a phosphorimager (Molecular Dynamics).

## 7.2. RESULTS

CHO-K1 cells transfected with a vector encoding the complete human prototype HER4 polypeptide were selected for amplified expression in media containing increasing concentrations of methionine sulfoximine as outlined in Section 7.1., et seq., *supra*. Expression of HER4 was evaluated using the immunoassay described in Section 7.1.3., *supra*. Several transfected CHO-K1 cell clones stably expressing HER4 were isolated. One particular clone, CHO/HER4 21-2, was selected in media supplemented with 250  $\mu$ M MSX, and expresses high levels of HER4. CHO/HER4 21-2 cells have been deposited with the ATCC.

Recombinant HER4 expressed in CHO/HER4 cells migrated with an apparent Mr of 180,000, slightly less than HER2, whereas the parental CHO cells showed no cross-reactive bands (FIG. 7A). In addition, a 130 kDa band was also detected in the CHO/HER4 cells, and presumably represents a degradation product of the 180 kDa mature protein. CHO/HER4 cells were used to identify ligand specific binding and autophosphorylation of the HER4 tyrosine kinase (see Section 9., et seq., *infra*).

## 8. EXAMPLE: ASSAY FOR DETECTING EGFR-FAMILY LIGANDS

### 8.1. CELL LINES

A panel of four recombinant cell lines, each expressing a single member of the human EGFR-family, were generated for use in the tyrosine kinase stimulatory assay described in Section 8.2., below. The cell line CHO/HER4 3 was generated as described in Section 7.1.2, *supra*.

CHO/HER2 cells (clone 1-2500) were selected to express high levels of recombinant human p185<sup>erbB2</sup> by dihydrofolate reductase-induced gene amplification in dhfr-deficient CHO cells. The HER2 expression

plasmid, cDNeu, was generated by insertion of a full length HER2 coding sequence into a modified pCDM8 (Invitrogen, San Diego, CA) expression vector (Seed and Aruffo, 1987, Proc. Natl. Acad. Sci. U.S.A. 84: 3365-69) in which an expression cassette from pSV2DHFR (containing the murine dhfr cDNA driven by the SV40 early promoter) has been inserted at the pCDM8 vector's unique BamHI site. This construct drives

6 HER2 expression from the CMV immediate-early promoter.

NRHER5 cells (Velu et al., 1987, Science 1408-10) were obtained from Dr. Hsing-Jian Kung (Case Western Reserve University, Cleveland, OH). This murine cell line was clonally isolated from NR6 cells infected with a retrovirus stock carrying the human EGFR, and was found to have approximately  $10^6$  human EGFRs per cell.

10 The cell line 293/HER3 was selected for high level expression of p180<sup>erbB3</sup>. The parental cell line, 293 human embryonic kidney cells, constitutively expresses adenovirus E1a and have low levels of EGFR expression. This line was established by cotransfection of linearized cHER3 (Plowman et al., 1990, Proc. Natl. Acad. Sci. U.S.A. 87: 4905-09) and pMC1neoPolyA (neomycin selectable marker with an Herpes simplex thymidine kinase promoter, Stratagene), with selection in DMEM/F12 media containing 500µg/ml G418.

## 8.2. TYROSINE KINASE STIMULATION ASSAY

Cells were plated in 6-well tissue culture plates (Falcon), and allowed to attach at 37°C for 18-24 hr. Prior to the assay, the cells were changed to serum-free media for at least 1 hour. Cell monolayers were then incubated with the amounts of ligand preparations indicated in Section 7.3., below for 5 min at 37°C. Cells were then washed with PBS and solubilized on ice with 0.5 ml PBSTDS containing phosphatase inhibitors (10 mM NaHPO<sub>4</sub>, 7.25, 150 mM NaCl, 1% Triton X-100, 0.5% deoxycholate, 0.1% SDS, 0.2% sodium azide, 1 mM NaF, 1 mM EGTA, 4 mM sodium orthovanadate, 1% aprotinin, 5 µg/ml leupeptin). Cell debris was removed by centrifugation (12000 x g, 15 min, 4°C) and the cleared supernatant reacted with 1 µg murine monoclonal antibody to phosphotyrosine (PY20, ICN Biochemicals, Cleveland, Ohio) for CHO/HER4 and 293/HER3 cells, or 1 µg murine monoclonal antibody to HER2 (Neu-Ab3, Oncogene Sciences) for CHO/HER2 cells, or 1 µg murine monoclonal antibody EGFR-1 to human EGFR (Amersham) for NRHER5 cells. Following a 1 hr incubation at 4°C, 30 µl of a 1:1 slurry (in PBSTDS) of anti-mouse IgG-agarose (for PY20 and Neu-Ab3 antibodies) or protein A-sepharose (for EGFR-R1 antibody) was added and the incubation was allowed to continue an additional 30 minutes. The beads were washed 3 times in PBSTDS and the complexes resolved by electrophoresis on reducing 7% SDS-polyacrylamide gels. The gels were transferred to nitrocellulose and blocked in TNET (10 mM Tris pH7.4, 75 mM NaCl, 0.1% Tween-20, 1 mM EDTA). PY20 antiphosphotyrosine antibody diluted 1:1000 in TNET was used as the primary antibody followed by <sup>125</sup>I-goat anti-mouse Ig F(ab')<sub>2</sub> diluted 1:500 in TNET. Blots were washed with TNET and exposed on a phosphorimager (Molecular Dynamics).

## 8.3. RESULTS

40 Several EGF-family member polypeptide and ligand preparations were tested for their ability to stimulate tyrosine phosphorylation of each of four EGFR-family receptors expressed in recombinant CHO cells using the tyrosine phosphorylation stimulation assay described in Section 8.2., above. The particular preparations tested for each of the four recombinant cell lines and the results obtained in the assay are tabulated below, and autoradiographs of some of these results are shown in FIG. 8.

TABLE III

PREPARATION	STIMULATION OF TYR PHOSPHORYLATION OF EGFR-FAMILY RECEPTORS			
	RECOMBINANT CELLS			
	CHO/HER4#3	CHO/HER2	NRHER5	293/HER3
EGF	-	-	+	-
AMPHIREGULIN	-	-	+	-
TGF- $\alpha$	-	-	+	-
HB-EGF	-	-	+	-
FRACTION 17*	+	-	-	-
FRACTION 14*	-	-	-	-

\* The identification of the HER4 tyrosine kinase stimulatory activity within the conditioned media of HepG2 cells and the isolation of these preparations is described in Section 8, *infra*.

The results indicate that EGF, AR, TGF- $\alpha$ , and HB-EGF, four related ligands which mediate their growth regulatory signals in part through interaction with EGFR, were able to stimulate tyrosine phosphorylation of EGFR expressed in recombinant NIH3T3 cells (for EGF, see FIG. 8C, lane 2), but not HER4, HER2, or HER3 expressed in recombinant CHO or 293 cells (FIG. 8A, B, D, lanes 2 and 3). Additionally, as discussed in more detail below, the assay identified a HepG2-derived preparation (fraction 17) as a HER4 ligand capable of specifically stimulating tyrosine phosphorylation of HER4 expressed in CHO/HER4 cells alone.

## 9. EXAMPLE: ISOLATION OF A HER4 LIGAND

### 9.1. MATERIALS AND METHODS

#### 9.1.1. CELL DIFFERENTIATION ASSAY

For the identification of ligands specific for HER2, HER3 or HER4, the receptor expression profile of MDA-MB-453 cells offers an excellent indicator for morphologic differentiation inducing activity. This cell line is known to express HER2 and HER3, but contains no detectable EGFR. The results of the semi-quantitative PCR assays (Table III) indicated high level expression of HER4 in MDA-MB-453 cells. In addition, cDNA encoding the prototype HER4 polypeptide of the invention was first isolated from this cell line (Section 6., *supra*).

MDA-MB-453 cells (7500/well) were grown in 50 ml DMEM supplemented with 5% FBS and 1x essential amino acids. Cells were allowed to adhere to 96-well plates for 24 hr. Samples were diluted in the above medium, added to the cell monolayer in 50 ml final volume, and the incubation continued for an additional 3 days. Cells were then examined by inverted light microscopy for morphologic changes.

#### 9.1.2. SOURCE CELLS

Serum free media from a panel of cultures human cancer cells were screened for growth regulatory activity on MDA-MB-453 cells. A human hepatocarcinoma cell line, HepG2, was identified as a source of a factor which induced dramatic morphologic differentiation of the MDA-MB-453 cells.

#### 9.1.3. PURIFICATION OF HER4 LIGAND

The cell differentiation assay described in Section 10.1.1., *supra*, was used throughout the purification procedure to monitor the column fractions that induce morphological changes in MDA-MB-453 cells. For large-scale production of conditioned medium, HepG2 cells were cultured in DMEM containing 10% fetal bovine serum using Nunc cell factories. At about 70% confluence, cells were washed then incubated with serum-free DMEM. Conditioned medium (HepG2-CM) was collected 3 days later, and fresh serum-free medium added to the cells. Two additional harvests of HepG2-CM were collected per cell factory. The medium was centrifuged and stored at -20°C in the presence of 500 mM PMSF.

Ten litres of HepG2-CM were concentrated 16-fold using an Amicon ultrafiltration unit (10,000 molecular weight cutoff membrane), and subjected to sequential precipitation with 20% and 60% ammonium sulfate. After centrifugation at 15,000 x g, the supernatant was extensively dialyzed against PBS and passed through a DEAE-sapharose (Pharmacia) column pre-equilibrated with PBS. The flow-through fraction was then applied onto a 4 ml heparin-acyrylic (Bio-Rad) column equilibrated with PBS. Differentiation inducing activity eluted from the heparin column between 0.4 and 0.8 M NaCl. Active heparin fractions were pooled, brought to 2.0 M ammonium sulfate, centrifuged at 12,000 x g for 5 min, and the resulting supernatant was loaded onto a phenyl-5PW column (8 x 75 mm, Waters). Bound proteins were eluted with a decreasing gradient from 2.0 M ammonium sulfate in 0.1 M Na<sub>2</sub>HPO<sub>4</sub>, pH 7.4 to 0.1 M Na<sub>2</sub>HPO<sub>4</sub>. Dialyzed fractions were assayed for tyrosine phosphorylation of MDA-MB-453 cells, essentially as described (Wen et al., 1992, Cell 69: 559-72), except PY20 was used as the primary antibody and horseradish peroxidase-conjugated goat F(ab')<sub>2</sub> anti-mouse Ig (Cappel) and chemiluminescence were used for detection. Phosphorylation signals were analyzed using the Molecular Dynamics personal densitometer.

## 9.2. RESULTS

Semi-purified HepG2-derived factor demonstrated a capacity to induce differentiation in MDA-MB-453 cells (FIG. 9). With reference to the micrographs shown in FIG. 9, untreated MDA-MB-453 cells are moderately adherent and show a rounded morphology (FIG. 9A). In contrast, the addition of semi-purified HepG2-derived factor induces these cells to display a noticeably flattened morphology with larger nuclei and increased cytoplasm (FIG. 9B and 9C). This HepG2-derived factor preparation also binds to heparin, a property which was utilized for purifying the activity.

On further purification, the HepG2-derived factor was found to elute from a phenyl hydrophobic interaction column at 1.0M ammonium sulfate (fractions 16 to 18). FIG. 9D shows the phenyl column elution profile. Tyrosine phosphorylation assays of the phenyl column fractions revealed that the same fractions found to induce differentiation of the human breast carcinoma cells are also able to stimulate tyrosine phosphorylation of a 185 K protein in MDA-MB-453 cells (FIG. 9E). In particular, fraction 16 induced a 4.5-fold increase in the phosphorylation signal compared to the baseline signal observed in unstimulated cells, as determined by densitometry analysis (FIG. 9F).

The phenyl fractions were also tested against the panel of cell lines which each overexpress a single member of the EGFR-family (Section 9.1., *supra*). Fraction 17 induced a significant and specific activation of the HER4 kinase (FIG. 8A, lane 4) without directly affecting the phosphorylation of HER2, EGFR, or HER3 (FIGS. 8B, 8C, and 8D, lane 4). Adjacent fraction 14 was used as a control and had no effect on the phosphorylation of any of the EGFR-family receptors (FIGS. 8A, B, C, D, lane 5). Further purification and analysis of the factor present in fraction 17 indicates that it is a glycoprotein of 40 to 45 kDa, approximately the same size as NDF and HRG. The HepG2-derived factor also has functional properties similar to NDF and HRG, inasmuch as it stimulates tyrosine phosphorylation of HER2/p185 in MDA-MB-453 cells, but not EGFR in NR5 cells, and induces morphologic differentiation of HER2 overexpressing human breast cancer cells.

Recently, several groups have reported the identification of specific ligands for HER2 (see Section 2., *supra*), including NDF and HRG- $\alpha$ . In contrast to these molecules, the HepG2-derived factor described herein failed to stimulate phosphorylation of HER2 in CHO/HER2 cells, but did stimulate phosphorylation of HER4 in CHO/HER4 cells. These findings are intriguing in view of the ability of the HepG2-derived factor to stimulate phosphorylation of MDA-MB-453 cells, a cell line known to overexpress HER2 and HER3 and the source from which HER4 was cloned. Since EGFR and HER2 have been shown to act synergistically, it is conceivable that HER4 may also interact with other EGFR-family members. In this connection, these results suggest that NDF may bind to HER4 in MDA-MB-453 cells resulting in the activation of HER2. The results described in Section 10., immediately below, provide evidence that NDF interacts directly with HER4, resulting in activation of HER2.

## 10. EXAMPLE: RECOMBINANT NDF-INDUCED, HER4 MEDIATED PHOSPHORYLATION OF HER2

Recombinant NDF was expressed in COS cells and tested for its activity on HER4 in an assay system essentially devoid of other known members of the EGFR-family, notably EGFR and HER2.

A full length rat NDF cDNA was isolated from normal rat kidney RNA and inserted into a cDM8-based expression vector to generate cNDF1.6. This construct was transiently expressed in COS cells, and conditioned cell supernatants were tested for NDF activity using the tyrosine kinase stimulation assay described in Section 8.2., *supra*. Supernatants from cNDF1.6 transfected cells upregulated tyrosine

phosphorylation in MDA-MB-453 cells relative to mock transfected COS media FIG. 10A. Phosphorylation peaked 10-15 minutes after addition on NDF.

The crude NDF supernatants were also tested for the ability to phosphorylate EGFR (NR5 cells), HER2 (CHO/HER2 1-2500 cells), and HER4 (CHO/HER4 21-2 cells). The NDF preparation had no effect on phosphorylation of EGFR, or HER2 containing cells, but induced a 2.4 to 4 fold increase in tyrosine phosphorylation of HER4 after 15 minutes incubation (see FIG. 10B). These findings provide preliminary evidence that NDF/HRG- $\alpha$  mediate their effects not through direct binding to HER2, but instead by means of a direct interaction with HER4. In cell lines expressing both HER2 and HER4, such as MDA-MB-453 cells and other breast carcinoma cells, binding of NDF to HER4 may stimulate HER2 either by heterodimer formation of these two related transmembrane receptors, or by intracellular crosstalk. Formal proof of the direct interaction between NDF and HER4 will require crosslinking of  $^{125}$ I-NDF to CHO/HER4 cells and a detailed analysis of its binding characteristics.

#### 11. EXAMPLE: CHROMOSOMAL MAPPING OF THE HER4 GENE

A HER4 cDNA probe corresponding to the 5' portion of the gene (nucleotide positions 34-1303) was used for *in situ* hybridization mapping of the HER4 gene. *In situ* hybridization to metaphase chromosomes from lymphocytes of two normal male donors was conducted using the HER4 probe labeled with  $^3$ H to a specific activity of  $2.6 \times 10^7$  cpm/ $\mu$ g as described (Marth et al, 1988, Proc. Natl. Acad. Sci. U.S.A. 83:7400-04). The final probe concentration was 0.05  $\mu$ g/ $\mu$ l of hybridization mixture. Slides were exposed for one month. Chromosomes were identified by Q banding.

#### 11.2 RESULTS

A total of 58 metaphase cells with autoradiographic grains were examined. Of the 124 hybridization sites scored, 38 (31%) were located on the distal portion of the long arm of chromosome 2 (FIG. 11). The greatest number of grains (21 grains) was located at band q33, with significant numbers of grains on bands q34 (10 grains) and q35 (7 grains). No significant hybridization on other human chromosomes was detected.

#### 12. MICROORGANISM AND CELL DEPOSITS

The following microorganisms and cell lines have been deposited with the American Type Culture Collection, and have been assigned the following accession numbers:

Microorganism	Plasmid	Accession Number
<i>Escherichia coli</i> SCS-1	pBSHER4Y	69 131
(containing the complete human HER4 coding sequence)		

Cell Lines	Accession Number
CHO/HER4 21-2	CRL 11205

The present invention is not to be limited in scope by the microorganisms and cell lines deposited or the embodiments disclosed herein, which are intended as single illustrations of one aspect of the invention, and any which are functionally equivalent are within the scope of the invention. Indeed, various modifications of the invention, in addition to those shown and described herein, will become apparent to those skilled in the art from the foregoing description. Such modifications are intended to fall within the scope of the appended claims. All base pair and amino acid residue numbers and sizes given for polynucleotides and polypeptides are approximate and used for the purpose of description.



SEQUENCE LISTING

6

(1) GENERAL INFORMATION:

10

(i) APPLICANT:

- (A) NAME: BRISTOL-MYERS SQUIBB COMPANY
- (B) STREET: 345 Park Avenue
- (C) CITY: New York
- (D) STATE: New York
- (E) COUNTRY: U.S.A.
- (F) POSTAL CODE (ZIP): 10154

15

(ii) TITLE OF INVENTION: HER4 HUMAN RECEPTOR TYROSINE KINASE

20

(iii) NUMBER OF SEQUENCES: 30

25

(iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.25

30

35

(2) INFORMATION FOR SEQ ID NO:1:

40

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5501 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

45

(ii) MOLECULE TYPE: DNA (genomic)

50

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 34..3961

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

	AATTGTCAGC	ACGGGATCTG	AGACTTCCAA	AAA	ATG	AAG	CCG	GCG	ACA	GGA	CTT	54	
					Met	Lys	Pro	Ala	Thr	Gly	Leu		
					1					5			
5	TGG	GTC	TGG	GTG	AGC	CTT	CTC	GTG	GCG	GCG	GGG	ACC	102
	Trp	Val	Trp	Val	Ser	Leu	Leu	Val	Ala	Ala	Gly	Thr	
			10					15				20	
	GAT	TCT	CAG	TCA	GTG	TGT	GCA	GGA	ACG	GAG	AAT	AAA	150
10	Asp	Ser	Gln	Ser	Val	Cys	Ala	Gly	Thr	Glu	Asn	Lys	
		25				30						35	
	TCT	GAC	CTG	GAA	CAG	CAG	TAC	CGA	GCC	TTG	CGC	AAG	198
	Ser	Asp	Leu	Glu	Gln	Gln	Tyr	Arg	Ala	Leu	Arg	Lys	
		40				45					50		
													55
15	TGT	GAG	GTT	GTG	ATG	GGC	AAC	CTG	GAG	ATA	ACC	AGC	246
	Cys	Glu	Val	Val	Met	Gly	Asn	Leu	Glu	Ile	Thr	Ser	
					60					65			
													70
	CGG	GAC	CTC	TCC	TTC	CTG	CGG	TCT	GTT	CGA	GAA	GTG	294
20	Arg	Asp	Leu	Ser	Phe	Leu	Arg	Ser	Val	Arg	Glu	Val	
				75					80				
													85
	TTA	GTG	GCT	CTT	AAT	CAG	TTT	CGT	TAC	CTG	CCT	CTG	342
	Leu	Val	Ala	Leu	Asn	Gln	Phe	Arg	Tyr	Leu	Pro	Leu	
			90					95					
													100
25	ATT	ATT	CGT	GGG	ACA	AAA	CTT	TAT	GAG	GAT	CGA	TAT	390
	Ile	Ile	Arg	Gly	Thr	Lys	Leu	Tyr	Glu	Asp	Arg	Tyr	
			105				110						
													115
	TTT	TTA	AAC	TAC	AGA	AAA	GAT	GGA	AAC	TTT	GGA	CTT	438
30	Phe	Leu	Asn	Tyr	Arg	Lys	Asp	Gly	Asn	Phe	Gly	Leu	
						125					130		
													135
	TTA	AAG	AAC	TTG	ACA	GAA	ATC	CTA	AAT	GGT	GGA	GTG	486
	Leu	Lys	Asn	Leu	Thr	Glu	Ile	Leu	Asn	Gly	Gly	Val	
					140					145			
													150
35	AAC	AAA	TTC	CTT	TGT	TAT	GCA	GAC	ACC	ATT	CAT	TGG	534
	Asn	Lys	Phe	Leu	Cys	Tyr	Ala	Asp	Thr	Ile	His	Trp	
				155					160				
													165
	CGG	AAC	CCA	TGG	CCT	TCC	AAC	TTG	ACT	CTT	GTG	TCA	582
40	Arg	Asn	Pro	Trp	Pro	Ser	Asn	Leu	Thr	Leu	Val	Ser	
			170					175					
													180
	TCA	GGA	TGT	GGA	CGT	TGC	CAT	AAG	TCC	TGT	ACT	GGC	630
	Ser	Gly	Cys	Gly	Arg	Cys	His	Lys	Ser	Cys	Thr	Gly	
			185				190						
													195
45	CCC	ACA	GAA	AAT	CAT	TGC	CAG	ACT	TTG	ACA	AGG	ACG	678
	Pro	Thr	Glu	Asn	His	Cys	Gln	Thr	Leu	Thr	Arg	Thr	
			200			205					210		
													215

50

55

	CAA TGT GAC GGC AGA TGC TAC GGA CCT TAC GTC AGT GAC TGC TGC CAT Gln Cys Asp Gly Arg Cys Tyr Gly Pro Tyr Val Ser Asp Cys Cys His 220 225 230	726
5	CGA GAA TGT GCT GGA GGC TGC TCA GGA CCT AAG GAC ACA GAC TGC TTT Arg Glu Cys Ala Gly Gly Cys Ser Gly Pro Lys Asp Thr Asp Cys Phe 235 240 245	774
10	GCC TGC ATG AAT TTC AAT GAC AGT GGA GCA TGT GTT ACT CAG TGT CCC Ala Cys Met Asn Phe Asn Asp Ser Gly Ala Cys Val Thr Gln Cys Pro 250 255 260	822
	CAA ACC TTT GTC TAC AAT CCA ACC ACC TTT CAA CTG GAG CAC AAT TTC Gln Thr Phe Val Tyr Asn Pro Thr Thr Phe Gln Leu Glu His Asn Phe 265 270 275	870
15	AAT GCA AAG TAC ACA TAT GGA GCA TTC TGT GTC AAG AAA TGT CCA CAT Asn Ala Lys Tyr Thr Tyr Gly Ala Phe Cys Val Lys Lys Cys Pro His 280 285 290 295	918
20	AAC TTT GTG GTA GAT TCC AGT TCT TGT GTG CGT GCC TGC CCT AGT TCC Asn Phe Val Val Asp Ser Ser Ser Cys Val Arg Ala Cys Pro Ser Ser 300 305 310	966
	AAG ATG GAA GTA GAA GAA AAT GGG ATT AAA ATG TGT AAA CCT TGC ACT Lys Met Glu Val Glu Glu Asn Gly Ile Lys Met Cys Lys Pro Cys Thr 315 320 325	1014
25	GAC ATT TGC CCA AAA GCT TGT GAT GGC ATT GGC ACA GGA TCA TTG ATG Asp Ile Cys Pro Lys Ala Cys Asp Gly Ile Gly Thr Gly Ser Leu Met 330 335 340	1062
30	TCA GCT CAG ACT GTG GAT TCC AGT AAC ATT GAC AAA TTC ATA AAC TGT Ser Ala Gln Thr Val Asp Ser Ser Asn Ile Asp Lys Phe Ile Asn Cys 345 350 355	1110
	ACC AAG ATC AAT GGG AAT TTG ATC TTT CTA GTC ACT GGT ATT CAT GGG Thr Lys Ile Asn Gly Asn Leu Ile Phe Leu Val Thr Gly Ile His Gly 360 365 370 375	1158
35	GAC CCT TAC AAT GCA ATT GAA GCC ATA GAC CCA GAG AAA CTG AAC GTC Asp Pro Tyr Asn Ala Ile Glu Ala Ile Asp Pro Glu Lys Leu Asn Val 380 385 390	1206
40	TTT CGG ACA GTC AGA GAG ATA ACA GGT TTC CTG AAC ATA CAG TCA TGG Phe Arg Thr Val Arg Glu Ile Thr Gly Phe Leu Asn Ile Gln Ser Trp 395 400 405	1254
	CCA CCA AAC ATG ACT GAC TTC AGT GTT TTT TCT AAC CTG GTG ACC ATT Pro Pro Asn Met Thr Asp Phe Ser Val Phe Ser Asn Leu Val Thr Ile 410 415 420	1302
45	GGT GGA AGA GTA CTC TAT AGT GGC CTG TCC TTG CTT ATC CTC AAG CAA Gly Gly Arg Val Leu Tyr Ser Gly Leu Ser Leu Leu Ile Leu Lys Gln 425 430 435	1350

50

55

	CAG GGC ATC ACC TCT CTA CAG TTC CAG TCC CTG AAG GAA ATC AGC GCA Gln Gly Ile Thr Ser Leu Gln Phe Gln Ser Leu Lys Glu Ile Ser Ala 440 445 450 455	1398
5	GGA AAC ATC TAT ATT ACT GAC AAC AGC AAC CTG TGT TAT TAT CAT ACC Gly Asn Ile Tyr Ile Thr Asp Asn Ser Asn Leu Cys Tyr Tyr His Thr 460 465 470	1446
10	ATT AAC TGG ACA ACA CTC TTC AGC ACA ATC AAC CAG AGA ATA GTA ATC Ile Asn Trp Thr Thr Leu Phe Ser Thr Ile Asn Gln Arg Ile Val Ile 475 480 485	1494
	CGG GAC AAC AGA AAA GCT GAA AAT TGT ACT GCT GAA GGA ATG GTG TGC Arg Asp Asn Arg Lys Ala Glu Asn Cys Thr Ala Glu Gly Met Val Cys 490 495 500	1542
15	AAC CAT CTG TGT TCC AGT GAT GGC TGT TGG GGA CCT GGG CCA GAC CAA Asn His Leu Cys Ser Ser Asp Gly Cys Trp Gly Pro Gly Pro Asp Gln 505 510 515	1590
20	TGT CTG TCG TGT CGC CGC TTC AGT AGA GGA AGG ATC TGC ATA GAG TCT Cys Leu Ser Cys Arg Phe Ser Arg Gly Arg Ile Cys Ile Glu Ser 520 525 530 535	1638
	TGT AAC CTC TAT GAT GGT GAA TTT CGG GAG TTT GAG AAT GGC TCC ATC Cys Asn Leu Tyr Asp Gly Glu Phe Arg Glu Phe Glu Asn Gly Ser Ile 540 545 550	1686
25	TGT GTG GAG TGT GAC CCC CAG TGT GAG AAG ATG GAA GAT GGC CTC CTC Cys Val Glu Cys Asp Pro Gln Cys Glu Lys Met Glu Asp Gly Leu Leu 555 560 565	1734
	ACA TGC CAT GGA CCG GGT CCT GAC AAC TGT ACA AAG TGC TCT CAT TTT Thr Cys His Gly Pro Gly Pro Asp Asn Cys Thr Lys Cys Ser His Phe 570 575 580	1782
30	AAA GAT GGC CCA AAC TGT GTG GAA AAA TGT CCA GAT GGC TTA CAG GGG Lys Asp Gly Pro Asn Cys Val Glu Lys Cys Pro Asp Gly Leu Gln Gly 585 590 595	1830
35	GCA AAC AGT TTC ATT TTC AAG TAT GCT GAT CCA GAT CGG GAG TGC CAC Ala Asn Ser Phe Ile Phe Lys Tyr Ala Asp Pro Asp Arg Glu Cys His 600 605 610 615	1878
	CCA TGC CAT CCA AAC TGC ACC CAA GGG TGT AAC GGT CCC ACT AGT CAT Pro Cys His Pro Asn Cys Thr Gln Gly Cys Asn Gly Pro Thr Ser His 620 625 630	1926
40	GAC TGC ATT TAC TAC CCA TGG ACG GGC CAT TCC ACT TTA CCA CAA CAT Asp Cys Ile Tyr Tyr Pro Trp Thr Gly His Ser Thr Leu Pro Gln His 635 640 645	1974
45	GCT AGA ACT CCC CTG ATT GCA GCT GGA GTA ATT GGT GGG CTC TTC ATT Ala Arg Thr Pro Leu Ile Ala Ala Gly Val Ile Gly Gly Leu Phe Ile 650 655 660	2022

50

55

	CTG GTC ATT GTG GGT CTG ACA TTT GCT GTT TAT GTT AGA AGG AAG AGC	2070
	Leu Val Ile Val Gly Leu Thr Phe Ala Val Tyr Val Arg Arg Lys Ser	
	665 670 675	
5	ATC AAA AAG AAA AGA GCC TTG AGA AGA TTC TTG GAA ACA GAG TTG GTG	2118
	Ile Lys Lys Lys Arg Ala Leu Arg Arg Phe Leu Glu Thr Glu Leu Val	
	680 685 690 695	
	GAA CCA TTA ACT CCC AGT GGC ACA GCA CCC AAT CAA GCT CAA CTT CGT	2166
	Glu Pro Leu Thr Pro Ser Gly Thr Ala Pro Asn Gln Ala Gln Leu Arg	
	700 705 710	
10	ATT TTG AAA GAA ACT GAG CTG AAG AGG GTA AAA GTC CTT GGC TCA GGT	2214
	Ile Leu Lys Glu Thr Glu Leu Lys Arg Val Lys Val Leu Gly Ser Gly	
	715 720 725	
15	GCT TTT GGA ACG GTT TAT AAA GGT ATT TGG GTA CCT GAA GGA GAA ACT	2262
	Ala Phe Gly Thr Val Tyr Lys Gly Ile Trp Val Pro Glu Gly Glu Thr	
	730 735 740	
	GTG AAG ATT CCT GTG GCT ATT AAG ATT CTT AAT GAG ACA ACT GGT CCC	2310
	Val Lys Ile Pro Val Ala Ile Lys Ile Leu Asn Glu Thr Thr Gly Pro	
	745 750 755	
20	AAG GCA AAT GTG GAG TTC ATG GAT GAA GCT CTG ATC ATG GCA AGT ATG	2358
	Lys Ala Asn Val Glu Phe Met Asp Glu Ala Leu Ile Met Ala Ser Met	
	760 765 770 775	
	GAT CAT CCA CAC CTA GTC CGG TTG CTG GGT GTG TGT CTG AGC CCA ACC	2406
	Asp His Pro His Leu Val Arg Leu Leu Gly Val Cys Leu Ser Pro Thr	
	780 785 790	
	ATC CAG CTG GTT ACT CAA CTT ATG CCC CAT GGC TGC CTG TTG GAG TAT	2454
	Ile Gln Leu Val Thr Gln Leu Met Pro His Gly Cys Leu Leu Glu Tyr	
	795 800 805	
30	GTC CAC GAG CAC AAG GAT AAC ATT GGA TCA CAA CTG CTG CTT AAC TGG	2502
	Val His Glu His Lys Asp Asn Ile Gly Ser Gln Leu Leu Leu Asn Trp	
	810 815 820	
	TGT GTC CAG ATA GCT AAG GGA ATG ATG TAC CTG GAA GAA AGA CGA CTC	2550
	Cys Val Gln Ile Ala Lys Gly Met Met Tyr Leu Glu Glu Arg Arg Leu	
	825 830 835	
35	GTT CAT CGG GAT TTG GCA GCC CGT AAT GTC TTA GTG AAA TCT CCA AAC	2598
	Val His Arg Asp Leu Ala Ala Arg Asn Val Leu Val Lys Ser Pro Asn	
	840 845 850 855	
	CAT GTG AAA ATC ACA GAT TTT GGG CTA GCC AGA CTC TTG GAA GGA GAT	2646
	His Val Lys Ile Thr Asp Phe Gly Leu Ala Arg Leu Leu Glu Gly Asp	
	860 865 870	
	GAA AAA GAG TAC AAT GCT GAT GGA GGA AAG ATG CCA ATT AAA TGG ATG	2694
	Glu Lys Glu Tyr Asn Ala Asp Gly Gly Lys Met Pro Ile Lys Trp Met	
	875 880 885	
45		
50		
55		

	GCT CTG GAG TGT ATA CAT TAC AGG AAA TTC ACC CAT CAG AGT GAC GTT	2742
	Ala Leu Glu Cys Ile His Tyr Arg Lys Phe Thr His Gln Ser Asp Val	
	890 895 900	
5	TGG AGC TAT GGA GTT ACT ATA TGG GAA CTG ATG ACC TTT GGA GGA AAA	2790
	Trp Ser Tyr Gly Val Thr Ile Trp Glu Leu Met Thr Phe Gly Gly Lys	
	905 91 915	
	CCC TAT GAT GGA ATT CCA ACG CGA GAA ATC CCT GAT TTA TTA GAG AAA	2838
	Pro Tyr Asp Gly Ile Pro Thr Arg Glu Ile Pro Asp Leu Leu Glu Lys	
10	920 925 930 935	
	GGA GAA CGT TTG CCT CAG CCT CCC ATC TGC ACT ATT GAC GTT TAC ATG	2886
	Gly Glu Arg Leu Pro Gln Pro Pro Ile Cys Thr Ile Asp Val Tyr Met	
	940 945 950	
15	GTC ATG GTC AAA TGT TGG ATG ATT GAT GCT GAC AGT AGA CCT AAA TTT	2934
	Val Met Val Lys Cys Trp Met Ile Asp Ala Asp Ser Arg Pro Lys Phe	
	955 960 965	
	AAG GAA CTG GCT GCT GAG TTT TCA AGG ATG GCT CGA GAC CCT CAA AGA	2982
	Lys Glu Leu Ala Ala Glu Phe Ser Arg Met Ala Arg Asp Pro Gln Arg	
20	970 975 980	
	TAC CTA GTT ATT CAG GGT GAT GAT CGT ATG AAG CTT CCC AGT CCA AAT	3030
	Tyr Leu Val Ile Gln Gly Asp Asp Arg Met Lys Leu Pro Ser Pro Asn	
	985 990 995	
25	GAC AGC AAG TTC TTT CAG AAT CTC TTG GAT GAA GAG GAT TTG GAA GAT	3078
	Asp Ser Lys Phe Phe Gln Asn Leu Leu Asp Glu Glu Asp Leu Glu Asp	
	1000 1005 1010 1015	
	ATG ATG GAT GCT GAG GAG TAC TTG GTC CCT CAG GCT TTC AAC ATC CCA	3126
	Met Met Asp Ala Glu Glu Tyr Leu Val Pro Gln Ala Phe Asn Ile Pro	
	1020 1025 1030	
30	CCT CCC ATC TAT ACT TCC AGA GCA AGA ATT GAC TCG AAT AGG AGT GAA	3174
	Pro Pro Ile Tyr Thr Ser Arg Ala Arg Ile Asp Ser Asn Arg Ser Glu	
	1035 1040 1045	
	ATT GGA CAC AGC CCT CCT CCT GCC TAC ACC CCC ATG TCA GGA AAC CAG	3222
	Ile Gly His Ser Pro Pro Pro Ala Tyr Thr Pro Met Ser Gly Asn Gln	
35	1050 1055 1060	
	TTT GTA TAC CGA GAT GGA GGT TTT GCT GCT GAA CAA GGA GTG TCT GTG	3270
	Phe Val Tyr Arg Asp Gly Gly Phe Ala Ala Glu Gln Gly Val Ser Val	
	1065 1070 1075	
40	CCC TAC AGA GCC CCA ACT AGC ACA ATT CCA GAA GCT CCT GTG GCA CAG	3318
	Pro Tyr Arg Ala Pro Thr Ser Thr Ile Pro Glu Ala Pro Val Ala Gln	
	1080 1085 1090 1095	
	GGT GCT ACT GCT GAG ATT TTT GAT GAC TCC TGC TGT AAT GGC ACC CTA	3366
	Gly Ala Thr Ala Glu Ile Phe Asp Asp Ser Cys Cys Asn Gly Thr Leu	
45	1100 1105 1110	

50

55

	CGC AAG CCA GTG GCA CCC CAT GTC CAA GAG GAC AGT AGC ACC CAG AGG Arg Lys Pro Val Ala Pro His Val Gln Glu Asp Ser Ser Thr Gln Arg 1115 1120 1125	3414
6	TAC AGT GCT GAC CCC ACC GTG TTT GCC CCA GAA CGG AGC CCA CGA GGA Tyr Ser Ala Asp Pro Thr Val Phe Ala Pro Glu Arg Ser Pro Arg Gly 1130 1135 1140	3462
10	GAG CTG GAT GAG GAA GGT TAC ATG ACT CCT ATG CGA GAC AAA CCC AAA Glu Leu Asp Glu Glu Gly Tyr Met Thr Pro Met Arg Asp Lys Pro Lys 1145 1150 1155	3510
	CAA GAA TAC CTG AAT CCA GTG GAG GAG AAC CCT TTT GTT TCT CGG AGA Gln Glu Tyr Leu Asn Pro Val Glu Glu Asn Pro Phe Val Ser Arg Arg 1160 1165 1170 1175	3558
15	AAA AAT GGA GAC CTT CAA GCA TTG GAT AAT CCC GAA TAT CAC AAT GCA Lys Asn Gly Asp Leu Gln Ala Leu Asp Asn Pro Glu Tyr His Asn Ala 1180 1185 1190	3606
20	TCC AAT GGT CCA CCC AAG GCC GAG GAT GAG TAT GTG AAT GAG CCA CTG Ser Asn Gly Pro Pro Lys Ala Glu Asp Glu Tyr Val Asn Glu Pro Leu 1195 1200 1205	3654
	TAC CTC AAC ACC TTT GCC AAC ACC TTG GGA AAA GCT GAG TAC CTG AAG Tyr Leu Asn Thr Phe Ala Asn Thr Leu Gly Lys Ala Glu Tyr Leu Lys 1210 1215 1220	3702
25	AAC AAC ATA CTG TCA ATG CCA GAG AAG GCC AAG AAA GCG TTT GAC AAC Asn Asn Ile Leu Ser Met Pro Glu Lys Ala Lys Lys Ala Phe Asp Asn 1225 1230 1235	3750
30	CCT GAC TAC TGG AAC CAC AGC CTG CCA CCT CGG AGC ACC CTT CAG CAC Pro Asp Tyr Trp Asn His Ser Leu Pro Pro Arg Ser Thr Leu Gln His 1240 1245 1250 1255	3798
	CCA GAC TAC CTG CAG GAG TAC AGC ACA AAA TAT TTT TAT AAA CAG AAT Pro Asp Tyr Leu Gln Glu Tyr Ser Thr Lys Tyr Phe Tyr Lys Gln Asn 1260 1265 1270	3846
35	GGG CGG ATC CGG CCT ATT GTG GCA GAG AAT CCT GAA TAC CTC TCT GAG Gly Arg Ile Arg Pro Ile Val Ala Glu Asn Pro Glu Tyr Leu Ser Glu 1275 1280 1285	3894
40	TTC TCC CTG AAG CCA GGC ACT GTG CTG CCG CCT CCA CCT TAC AGA CAC Phe Ser Leu Lys Pro Gly Thr Val Leu Pro Pro Pro Pro Tyr Arg His 1290 1295 1300	3942
	CGG AAT ACT GTG GTG TAAGCTCAGT TGTGGTTTTT TAGGTGGAGA GACACACCTG Arg Asn Thr Val Val 1305	3997
45	CTCCAATTTT CCCACCCCCC TCTCTTCTC TGGTGGTCTT CCTTCTACCC CAAGGCCAGT	4057
	AGTTTTGACA CTTCACAGTG GAAGATACAG AGATGCAATG ATAGTTATGT GCTTACCTAA	4117
	CTTGAACATT AGAGGGAAAG ACTGAAAGAG AAAGATAGGA GGAACCACAA TGTTTCTTCA	4177

50

55

TTTCTCTGCA TGGGTTGGTC AGGAGAATGA AACAGCTAGA GAAGGACCAG AAAATGTAAG 4237  
 GCAATGCTGC CTAATATCAA ACTAGCTGTC ACTTTTTTTC TTTTCTTTT TCTTCTTTG 4297  
 5 TTTCTTTCTT CCTCTTCTTT TTTTTTTTTT TTTTAAAGCA GATGGTTGAA ACACCCATGC 4357  
 TATCTGTTCC TATCTGCAGG AACTGATGTG TGCATATTTA GCATCCCTGG AAATCATAAT 4417  
 AAAGTTTCCA TTAGAACAAA AGAATAACAT TTTCTATAAC ATATGATAGT GTCTGAAATT 4477  
 10 GAGAATCCAG TTTCTTTCCC CAGCAGTTTC TGTCTAGCA AGTAAGAATG GCCAACTCAA 4537  
 CTTTCATAAT TTAAAAATCT CCATTAAAGT TATAACTAGT AATTATGTTT TCAACACTTT 4597  
 TTGGTTTTTT TCATTTTGTT TTGCTCTGAC CGATTCTTTT ATATTGCTC CCCTATTTTT 4657  
 15 GGCTTTAATT TCTAATTGCA AAGATGTTT CATCAAAGCT TCTTCACAGA ATTTAAGCAA 4717  
 GAAATATTTT AATATAGTGA AATGGCCACT ACTTTAAGTA TACAATCTTT AAAATAAGAA 4777  
 AGGGAGGCTA ATATTTTTC TGTATCAAA TTATCTTCAC CCTCATCCTT TACATTTTTT 4837  
 AACATTTTTT TTTCTCCATA AATGACACTA CTTGATAGGC CGTTGGTTGT CTGAAGAGTA 4897  
 20 GAAGGGAAC TAAGAGACAG TTCTCTGTGG TTCAGGAAAA CTAATGATAC TTTCAGGGGT 4957  
 GGCCCAATGA GGAATCCAT TGAAGTGAA GAAACACACT GGATTGGGTA TGTCTACCTG 5017  
 GCAGATACTC AGAAATGTAG TTTGCACITA AGCTGTAATT TTATTGTTC TTTTCTGAA 5077  
 25 CTCCATTTTG GATTTTGAAT CAAGCAATAT GGAAGCAACC AGCAAATTAA CTAATTTAAG 5137  
 TACATTTTTA AAAAAGAGC TAAGATAAAG ACTGTGGAAG TGCCAAACCA AGCAAATTAG 5197  
 GAACCTTGCA ACGGTATCCA GGGACTATGA TGAGAGGCCA GCACATTATC TTCATATGTC 5257  
 30 ACCTTTGCTA CGCAAGGAAA TTTGTTCACT TCGTATACTT CGTAAGAAGG AATGCGAGTA 5317  
 AGGATTGGCT TGAATTCCAT GGAATTTCTA GTATGAGACT ATTTATATGA AGTAGAAGGT 5377  
 AACTCTTTGC ACATAAATTG GTATAATAAA AAGAAAAACA CAAACATTCA AAGCTTAGGG 5437  
 35 ATAGGTCCTT GGGTCAAAAG TTGTAAATAA ATGTGAAACA TCTTCTCAA AAAAAAAAAA 5497  
 AAAA 5501

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1308 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:



Met Lys Pro Ala Thr Gly Leu Trp Val Trp Val Ser Leu Leu Val Ala  
 1 5 10 15  
 6 Ala Gly Thr Val Gln Pro Ser Asp Ser Gln Ser Val Cys Ala Gly Thr  
 20 25 30  
 Glu Asn Lys Leu Ser Ser Leu Ser Asp Leu Glu Gln Gln Tyr Arg Ala  
 35 40 45  
 10 Leu Arg Lys Tyr Tyr Glu Asn Cys Glu Val Val Met Gly Asn Leu Glu  
 50 55 60  
 Ile Thr Ser Ile Glu His Asn Arg Asp Leu Ser Phe Leu Arg Ser Val  
 65 70 75 80  
 15 Arg Glu Val Thr Gly Tyr Val Leu Val Ala Leu Asn Gln Phe Arg Tyr  
 85 90 95  
 Leu Pro Leu Glu Asn Leu Arg Ile Ile Arg Gly Thr Lys Leu Tyr Glu  
 100 105 110  
 20 Asp Arg Tyr Ala Leu Ala Ile Phe Leu Asn Tyr Arg Lys Asp Gly Asn  
 115 120 125  
 Phe Gly Leu Gln Glu Leu Gly Leu Lys Asn Leu Thr Glu Ile Leu Asn  
 130 135 140  
 25 Gly Gly Val Tyr Val Asp Gln Asn Lys Phe Leu Cys Tyr Ala Asp Thr  
 145 150 155 160  
 Ile His Trp Gln Asp Ile Val Arg Asn Pro Trp Pro Ser Asn Leu Thr  
 165 170 175  
 30 Leu Val Ser Thr Asn Gly Ser Ser Gly Cys Gly Arg Cys His Lys Ser  
 180 185 190  
 Cys Thr Gly Arg Cys Trp Gly Pro Thr Glu Asn His Cys Gln Thr Leu  
 195 200 205  
 35 Thr Arg Thr Val Cys Ala Glu Gln Cys Asp Gly Arg Cys Tyr Gly Pro  
 210 215 220  
 Tyr Val Ser Asp Cys Cys His Arg Glu Cys Ala Gly Gly Cys Ser Gly  
 225 230 235 240  
 Pro Lys Asp Thr Asp Cys Phe Ala Cys Met Asn Phe Asn Asp Ser Gly  
 245 250 255  
 45 Ala Cys Val Thr Gln Cys Pro Gln Thr Phe Val Tyr Asn Pro Thr Thr  
 260 265 270  
 Phe Gln Leu Glu His Asn Phe Asn Ala Lys Tyr Thr Tyr Gly Ala Phe  
 275 280 285  
 50 Cys Val Lys Lys Cys Pro His Asn Phe Val Val Asp Ser Ser Ser Cys  
 290 295 300

55

Val Arg Ala Cys Pro Ser Ser Lys Met Glu Val Glu Glu Asn Gly Ile  
 305 310 315 320  
 5 Lys Met Cys Lys Pr Cys Thr Asp Ile Cys Pro Lys Ala Cys Asp Gly  
 325 330 335  
 Ile Gly Thr Gly Ser Leu Met Ser Ala Gln Thr Val Asp Ser Ser Asn  
 340 345 350  
 10 Ile Asp Lys Phe Ile Asn Cys Thr Lys Ile Asn Gly Asn Leu Ile Phe  
 355 360 365  
 Leu Val Thr Gly Ile His Gly Asp Pro Tyr Asn Ala Ile Glu Ala Ile  
 370 375 380  
 15 Asp Pro Glu Lys Leu Asn Val Phe Arg Thr Val Arg Glu Ile Thr Gly  
 385 390 395 400  
 Phe Leu Asn Ile Gln Ser Trp Pro Pro Asn Met Thr Asp Phe Ser Val  
 405 410 415  
 20 Phe Ser Asn Leu Val Thr Ile Gly Gly Arg Val Leu Tyr Ser Gly Leu  
 420 425 430  
 Ser Leu Leu Ile Leu Lys Gln Gln Gly Ile Thr Ser Leu Gln Phe Gln  
 435 440 445  
 25 Ser Leu Lys Glu Ile Ser Ala Gly Asn Ile Tyr Ile Thr Asp Asn Ser  
 450 455 460  
 Asn Leu Cys Tyr Tyr His Thr Ile Asn Trp Thr Thr Leu Phe Ser Thr  
 465 470 475 480  
 30 Ile Asn Gln Arg Ile Val Ile Arg Asp Asn Arg Lys Ala Glu Asn Cys  
 485 490 495  
 Thr Ala Glu Gly Met Val Cys Asn His Leu Cys Ser Ser Asp Gly Cys  
 500 505 510  
 35 Trp Gly Pro Gly Pro Asp Gln Cys Leu Ser Cys Arg Arg Phe Ser Arg  
 515 520 525  
 Gly Arg Ile Cys Ile Glu Ser Cys Asn Leu Tyr Asp Gly Glu Phe Arg  
 530 535 540  
 40 Glu Phe Glu Asn Gly Ser Ile Cys Val Glu Cys Asp Pro Gln Cys Glu  
 545 550 555 560  
 Lys Met Glu Asp Gly Leu Leu Thr Cys His Gly Pro Gly Pro Asp Asn  
 565 570 575  
 45 Cys Thr Lys Cys Ser His Phe Lys Asp Gly Pro Asn Cys Val Glu Lys  
 580 585 590  
 50 Cys Pro Asp Gly Leu Gln Gly Ala Asn Ser Phe Ile Phe Lys Tyr Ala  
 595 600 605  
 55

Asp Pro Asp Arg Glu Cys His Pro Cys His Pro Asn Cys Thr Gln Gly  
 610 615 620  
 Cys Asn Gly Pro Thr Ser His Asp Cys Il Tyr Tyr Pr Trp Thr Gly  
 625 630 635 640  
 His Ser Thr Leu Pro Gln His Ala Arg Thr Pro Leu Ile Ala Ala Gly  
 645 650 655  
 Val Ile Gly Gly Leu Phe Ile Leu Val Ile Val Gly Leu Thr Phe Ala  
 660 665 670  
 Val Tyr Val Arg Arg Lys Ser Ile Lys Lys Lys Arg Ala Leu Arg Arg  
 675 680 685  
 Phe Leu Glu Thr Glu Leu Val Glu Pro Leu Thr Pro Ser Gly Thr Ala  
 690 695 700  
 Pro Asn Gln Ala Gln Leu Arg Ile Leu Lys Glu Thr Glu Leu Lys Arg  
 705 710 715 720  
 Val Lys Val Leu Gly Ser Gly Ala Phe Gly Thr Val Tyr Lys Gly Ile  
 725 730 735  
 Trp Val Pro Glu Gly Glu Thr Val Lys Ile Pro Val Ala Ile Lys Ile  
 740 745 750  
 Leu Asn Glu Thr Thr Gly Pro Lys Ala Asn Val Glu Phe Met Asp Glu  
 755 760 765  
 Ala Leu Ile Met Ala Ser Met Asp His Pro His Leu Val Arg Leu Leu  
 770 775 780  
 Gly Val Cys Leu Ser Pro Thr Ile Gln Leu Val Thr Gln Leu Met Pro  
 785 790 795 800  
 His Gly Cys Leu Leu Glu Tyr Val His Glu His Lys Asp Asn Ile Gly  
 805 810 815  
 Ser Gln Leu Leu Leu Asn Trp Cys Val Gln Ile Ala Lys Gly Met Met  
 820 825 830  
 Tyr Leu Glu Glu Arg Arg Leu Val His Arg Asp Leu Ala Ala Arg Asn  
 835 840 845  
 Val Leu Val Lys Ser Pro Asn His Val Lys Ile Thr Asp Phe Gly Leu  
 850 855 860  
 Ala Arg Leu Leu Glu Gly Asp Glu Lys Glu Tyr Asn Ala Asp Gly Gly  
 865 870 875 880  
 Lys Met Pro Ile Lys Trp Met Ala Leu Glu Cys Ile His Tyr Arg Lys  
 885 890 895  
 Phe Thr His Gln Ser Asp Val Trp Ser Tyr Gly Val Thr Ile Trp Glu  
 900 905 910

55

Leu Met Thr Phe Gly Gly Lys Pro Tyr Asp Gly Ile Pro Thr Arg Glu  
 915 920 925  
 Ile Pro Asp Leu Leu Glu Lys Gly Glu Arg Leu Pro Gln Pro Pro Ile  
 5 930 935 940  
 Cys Thr Ile Asp Val Tyr Met Val Met Val Lys Cys Trp Met Ile Asp  
 945 950 955 960  
 Ala Asp Ser Arg Pro Lys Phe Lys Glu Leu Ala Ala Glu Phe Ser Arg  
 10 965 970 975  
 Met Ala Arg Asp Pro Gln Arg Tyr Leu Val Ile Gln Gly Asp Asp Arg  
 980 985 990  
 Met Lys Leu Pro Ser Pro Asn Asp Ser Lys Phe Phe Gln Asn Leu Leu  
 15 995 1000 1005  
 Asp Glu Glu Asp Leu Glu Asp Met Met Asp Ala Glu Glu Tyr Leu Val  
 1010 1015 1020  
 Pro Gln Ala Phe Asn Ile Pro Pro Pro Ile Tyr Thr Ser Arg Ala Arg  
 20 1025 1030 1035 1040  
 Ile Asp Ser Asn Arg Ser Glu Ile Gly His Ser Pro Pro Pro Ala Tyr  
 1045 1050 1055  
 Thr Pro Met Ser Gly Asn Gln Phe Val Tyr Arg Asp Gly Gly Phe Ala  
 25 1060 1065 1070  
 Ala Glu Gln Gly Val Ser Val Pro Tyr Arg Ala Pro Thr Ser Thr Ile  
 1075 1080 1085  
 Pro Glu Ala Pro Val Ala Gln Gly Ala Thr Ala Glu Ile Phe Asp Asp  
 30 1090 1095 1100  
 Ser Cys Cys Asn Gly Thr Leu Arg Lys Pro Val Ala Pro His Val Gln  
 1105 1110 1115 1120  
 Glu Asp Ser Ser Thr Gln Arg Tyr Ser Ala Asp Pro Thr Val Phe Ala  
 35 1125 1130 1135  
 Pro Glu Arg Ser Pro Arg Gly Glu Leu Asp Glu Glu Gly Tyr Met Thr  
 1140 1145 1150  
 Pro Met Arg Asp Lys Pro Lys Gln Glu Tyr Leu Asn Pro Val Glu Glu  
 1155 1160 1165  
 Asn Pro Phe Val Ser Arg Arg Lys Asn Gly Asp Leu Gln Ala Leu Asp  
 45 1170 1175 1180  
 Asn Pro Glu Tyr His Asn Ala Ser Asn Gly Pro Pro Lys Ala Glu Asp  
 1185 1190 1195 1200  
 Glu Tyr Val Asn Glu Pro Leu Tyr Leu Asn Thr Phe Ala Asn Thr Leu  
 50 1205 1210 1215

55

Gly Lys Ala Glu Tyr Leu Lys Asn Asn Ile Leu Ser Met Pro Glu Lys  
 1220 1225 1230  
 6 Ala Lys Lys Ala Phe Asp Asn Pro Asp Tyr Trp Asn His Ser Leu Pro  
 1235 1240 1245  
 Pro Arg Ser Thr Leu Gln His Pro Asp Tyr Leu Gln Glu Tyr Ser Thr  
 1250 1255 1260  
 10 Lys Tyr Phe Tyr Lys Gln Asn Gly Arg Ile Arg Pro Ile Val Ala Glu  
 1265 1270 1275 1280  
 Asn Pro Glu Tyr Leu Ser Glu Phe Ser Leu Lys Pro Gly Thr Val Leu  
 1285 1290 1295  
 15 Pro Pro Pro Pro Tyr Arg His Arg Asn Thr Val Val  
 1300 1305

## (2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 5555 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 34..3210

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

	AATTGTCAGC ACGGGATCTG AGACTTCCAA AAA ATG AAG CCG GCG ACA GGA CTT	54
	Met Lys Pro Ala Thr Gly Leu	
	1 5	
35	TGG GTC TGG GTG AGC CTT CTC GTG GCG GCG GGG ACC GTC CAG CCC AGC	102
	Trp Val Trp Val Ser Leu Leu Val Ala Ala Gly Thr Val Gln Pro Ser	
	10 15 20	
	GAT TCT CAG TCA GTG TGT GCA GGA ACG GAG AAT AAA CTG AGC TCT CTC	150
40	Asp Ser Gln Ser Val Cys Ala Gly Thr Glu Asn Lys Leu Ser Ser Leu	
	25 30 35	
	TCT GAC CTG GAA CAG CAG TAC CGA GCC TTG CGC AAG TAC TAT GAA AAC	198
	Ser Asp Leu Glu Gln Gln Tyr Arg Ala Leu Arg Lys Tyr Tyr Glu Asn	
	40 45 50 55	
45	TGT GAG GTT GTC ATG GGC AAC CTG GAG ATA ACC AGC ATT GAG CAC AAC	246
	Cys Glu Val Val Met Gly Asn Leu Glu Ile Thr Ser Ile Glu His Asn	
	60 65 70	

50

55

	CGG GAC CTC TCC TTC CTG CGG TCT GTT CGA GAA GTC ACA GGC TAC GTG	294
	Arg Asp Leu Ser Phe Leu Arg Ser Val Arg Glu Val Thr Gly Tyr Val	
	75 80 85	
5	TTA GTG GCT CTT AAT CAG TTT CGT TAC CTG CCT CTG GAG AAT TTA CGC	342
	Leu Val Ala Leu Asn Gln Phe Arg Tyr Leu Pro Leu Glu Asn Leu Arg	
	90 95 100	
	ATT ATT CGT GGG ACA AAA CTT TAT GAG GAT CGA TAT GCC TTG GCA ATA	390
10	Ile Ile Arg Gly Thr Lys Leu Tyr Glu Asp Arg Tyr Ala Leu Ala Ile	
	105 110 115	
	TTT TTA AAC TAC AGA AAA GAT GGA AAC TTT GGA CTT CAA GAA CTT GGA	438
	Phe Leu Asn Tyr Arg Lys Asp Gly Asn Phe Gly Leu Gln Glu Leu Gly	
	120 125 130 135	
15	TTA AAG AAC TTG ACA GAA ATC CTA AAT GGT GGA GTC TAT GTA GAC CAG	486
	Leu Lys Asn Leu Thr Glu Ile Leu Asn Gly Gly Val Tyr Val Asp Gln	
	140 145 150	
	AAC AAA TTC CTT TGT TAT GCA GAC ACC ATT CAT TGG CAA GAT ATT GTT	534
20	Asn Lys Phe Leu Cys Tyr Ala Asp Thr Ile His Trp Gln Asp Ile Val	
	155 160 165	
	CGG AAC CCA TGG CCT TCC AAC TTG ACT CTT GTG TCA ACA AAT GGT AGT	582
	Arg Asn Pro Trp Pro Ser Asn Leu Thr Leu Val Ser Thr Asn Gly Ser	
	170 175 180	
25	TCA GGA TGT GGA CGT TGC CAT AAG TCC TGT ACT GGC CGT TGC TGG GGA	630
	Ser Gly Cys Gly Arg Cys His Lys Ser Cys Thr Gly Arg Cys Trp Gly	
	185 190 195	
	CCC ACA GAA AAT CAT TGC CAG ACT TTG ACA AGG ACG GTG TGT GCA GAA	678
30	Pro Thr Glu Asn His Cys Gln Thr Leu Thr Arg Thr Val Cys Ala Glu	
	200 205 210 215	
	CAA TGT GAC GGC AGA TGC TAC GGA CCT TAC GTC AGT GAC TGC TGC CAT	726
	Gln Cys Asp Gly Arg Cys Tyr Gly Pro Tyr Val Ser Asp Cys Cys His	
	220 225 230	
35	CGA GAA TGT GCT GGA GGC TGC TCA GGA CCT AAG GAC ACA GAC TGC TTT	774
	Arg Glu Cys Ala Gly Gly Cys Ser Gly Pro Lys Asp Thr Asp Cys Phe	
	235 240 245	
	GCC TGC ATG AAT TTC AAT GAC AGT GGA GCA TGT GTT ACT CAG TGT CCC	822
40	Ala Cys Met Asn Phe Asn Asp Ser Gly Ala Cys Val Thr Gln Cys Pro	
	250 255 260	
	CAA ACC TTT GTC TAC AAT CCA ACC ACC TTT CAA CTG GAG CAC AAT TTC	870
	Gln Thr Phe Val Tyr Asn Pro Thr Thr Phe Gln Leu Glu His Asn Phe	
	265 270 275	
45	AAT GCA AAG TAC ACA TAT GGA GCA TTC TGT GTC AAG AAA TGT CCA CAT	918
	Asn Ala Lys Tyr Thr Tyr Gly Ala Phe Cys Val Lys Lys Cys Pro His	
	280 285 290 295	

50

55

	AAC TTT GTG GTA GAT TCC AGT TCT TGT GTG CGT GCC TGC CCT AGT TCC	966
	Asn Phe Val Val Asp Ser Ser Ser Cys Val Arg Ala Cys Pro Ser Ser	
	300 305 310	
6	AAG ATG GAA GTA GAA GAA AAT GGG ATT AAA ATG TGT AAA CCT TGC ACT	1014
	Lys Met Glu Val Glu Glu Asn Gly Ile Lys Met Cys Lys Pro Cys Thr	
	315 320 325	
	GAC ATT TGC CCA AAA GCT TGT GAT GGC ATT GGC ACA GGA TCA TTG ATG	1062
	Asp Ile Cys Pro Lys Ala Cys Asp Gly Ile Gly Thr Gly Ser Leu Met	
	330 335 340	
10	TCA GCT CAG ACT GTG GAT TCC AGT AAC ATT GAC AAA TTC ATA AAC TGT	1110
	Ser Ala Gln Thr Val Asp Ser Ser Asn Ile Asp Lys Phe Ile Asn Cys	
	345 350 355	
15	ACC AAG ATC AAT GGG AAT TTG ATC TTT CTA GTC ACT GGT ATT CAT GGG	1158
	Thr Lys Ile Asn Gly Asn Leu Ile Phe Leu Val Thr Gly Ile His Gly	
	360 365 370 375	
	GAC CCT TAC AAT GCA ATT GAA GCC ATA GAC CCA GAG AAA CTG AAC GTC	1206
	Asp Pro Tyr Asn Ala Ile Glu Ala Ile Asp Pro Glu Lys Leu Asn Val	
	380 385 390	
20	TTT CGG ACA GTC AGA GAG ATA ACA GGT TTC CTG AAC ATA CAG TCA TGG	1254
	Phe Arg Thr Val Arg Glu Ile Thr Gly Phe Leu Asn Ile Gln Ser Trp	
	395 400 405	
25	CCA CCA AAC ATG ACT GAC TTC AGT GTT TTT TCT AAC CTG GTG ACC ATT	1302
	Pro Pro Asn Met Thr Asp Phe Ser Val Phe Ser Asn Leu Val Thr Ile	
	410 415 420	
	GGT GGA AGA GTA CTC TAT AGT GGC CTG TCC TTG CTT ATC CTC AAG CAA	1350
	Gly Gly Arg Val Leu Tyr Ser Gly Leu Ser Leu Leu Ile Leu Lys Gln	
	425 430 435	
30	CAG GGC ATC ACC TCT CTA CAG TTC CAG TCC CTG AAG GAA ATC AGC GCA	1398
	Gln Gly Ile Thr Ser Leu Gln Phe Gln Ser Leu Lys Glu Ile Ser Ala	
	440 445 450 455	
	GGA AAC ATC TAT ATT ACT GAC AAC AGC AAC CTG TGT TAT TAT CAT ACC	1446
	Gly Asn Ile Tyr Ile Thr Asp Asn Ser Asn Leu Cys Tyr Tyr His Thr	
	460 465 470	
	ATT AAC TGG ACA ACA CTC TTC AGC ACA ATC AAC CAG AGA ATA GTA ATC	1494
	Ile Asn Trp Thr Thr Leu Phe Ser Thr Ile Asn Gln Arg Ile Val Ile	
	475 480 485	
40	CGG GAC AAC AGA AAA GCT GAA AAT TGT ACT GCT GAA GGA ATG GTG TGC	1542
	Arg Asp Asn Arg Lys Ala Glu Asn Cys Thr Ala Glu Gly Met Val Cys	
	490 495 500	
	AAC CAT CTG TGT TCC AGT GAT GGC TGT TGG GGA CCT GGG CCA GAC CAA	1590
	Asn His Leu Cys Ser Ser Asp Gly Cys Trp Gly Pro Gly Pro Asp Gln	
	505 510 515	
45		
50		
55		

	TGT	CTG	TCG	TGT	CGC	CGC	TTC	AGT	AGA	GGA	AGG	ATC	TGC	ATA	GAG	TCT	1638
	Cys	Leu	Ser	Cys	Arg	Arg	Phe	Ser	Arg	Gly	Arg	Ile	Cys	Ile	Glu	Ser	
	520				525					530					535		
5	TGT	AAC	CTC	TAT	GAT	GGT	GAA	TTT	CGG	GAG	TTT	GAG	AAT	GGC	TCC	ATC	1686
	Cys	Asn	Leu	Tyr	Asp	Gly	Glu	Phe	Arg	Glu	Phe	Glu	Asn	Gly	Ser	Ile	
				540						545					550		
	TGT	GTG	GAG	TGT	GAC	CCC	CAG	TGT	GAG	AAG	ATG	GAA	GAT	GGC	CTC	CTC	1734
	Cys	Val	Glu	Cys	Asp	Pro	Gln	Cys	Glu	Lys	Met	Glu	Asp	Gly	Leu	Leu	
				555					560					565			
10	ACA	TGC	CAT	GGA	CCG	GGT	CCT	GAC	AAC	TGT	ACA	AAG	TGC	TCT	CAT	TTT	1782
	Thr	Cys	His	Gly	Pro	Gly	Pro	Asp	Asn	Cys	Thr	Lys	Cys	Ser	His	Phe	
				570				575					580				
15	AAA	GAT	GGC	CCA	AAC	TGT	GTG	GAA	AAA	TGT	CCA	GAT	GGC	TTA	CAG	GGG	1830
	Lys	Asp	Gly	Pro	Asn	Cys	Val	Glu	Lys	Cys	Pro	Asp	Gly	Leu	Gln	Gly	
		585					590					595					
	GCA	AAC	AGT	TTC	ATT	TTC	AAG	TAT	GCT	GAT	CCA	GAT	CGG	GAG	TGC	CAC	1878
	Ala	Asn	Ser	Phe	Ile	Phe	Lys	Tyr	Ala	Asp	Pro	Asp	Arg	Glu	Cys	His	
	600				605					610					615		
20	CCA	TGC	CAT	CCA	AAC	TGC	ACC	CAA	GGG	TGT	AAC	GGT	CCC	ACT	AGT	CAT	1926
	Pro	Cys	His	Pro	Asn	Cys	Thr	Gln	Gly	Cys	Asn	Gly	Pro	Thr	Ser	His	
				620					625						630		
	GAC	TGC	ATT	TAC	TAC	CCA	TGG	ACG	GGC	CAT	TCC	ACT	TTA	CCA	CAA	CAT	1974
25	Asp	Cys	Ile	Tyr	Tyr	Pro	Trp	Thr	Gly	His	Ser	Thr	Leu	Pro	Gln	His	
				635					640						645		
	GCT	AGA	ACT	CCC	CTG	ATT	GCA	GCT	GGA	GTA	ATT	GGT	GGG	CTC	TTC	ATT	2022
	Ala	Arg	Thr	Pro	Leu	Ile	Ala	Gly	Val	Ile	Gly	Gly	Leu	Phe	Ile		
				650				655					660				
30	CTG	GTC	ATT	GTG	GGT	CTG	ACA	TTT	GCT	GTT	TAT	GTT	AGA	AGG	AAG	AGC	2070
	Leu	Val	Ile	Val	Gly	Leu	Thr	Phe	Ala	Val	Tyr	Val	Arg	Arg	Lys	Ser	
				665			670					675					
	ATC	AAA	AAG	AAA	AGA	GCC	TTG	AGA	AGA	TTC	TTG	GAA	ACA	GAG	TTG	GTG	2118
35	Ile	Lys	Lys	Lys	Arg	Ala	Leu	Arg	Arg	Phe	Leu	Glu	Thr	Glu	Leu	Val	
		680				685				690					695		
	GAA	CCA	TTA	ACT	CCC	AGT	GGC	ACA	GCA	CCC	AAT	CAA	GCT	CAA	CTT	CGT	2166
	Glu	Pro	Leu	Thr	Pro	Ser	Gly	Thr	Ala	Pro	Asn	Gln	Ala	Gln	Leu	Arg	
				700						705					710		
40	ATT	TTG	AAA	GAA	ACT	GAG	CTG	AAG	AGG	GTA	AAA	GTC	CTT	GGC	TCA	GGT	2214
	Ile	Leu	Lys	Glu	Thr	Glu	Leu	Lys	Arg	Val	Lys	Val	Leu	Gly	Ser	Gly	
				715				720						725			
	GCT	TTT	GGA	ACG	GTT	TAT	AAA	GGT	ATT	TGG	GTA	CCT	GAA	GGA	GAA	ACT	2262
45	Ala	Phe	Gly	Thr	Val	Tyr	Lys	Gly	Ile	Trp	Val	Pro	Glu	Gly	Glu	Thr	
			730					735					740				
50																	
55																	



	GTG AAG ATT CCT GTG GCT ATT AAG ATT CTT AAT GAG ACA ACT GGT CCC	2310
	Val Lys Ile Pro Val Ala Ile Lys Ile Leu Asn Glu Thr Thr Gly Pro	
	745 750 755	
6	AAG GCA AAT GTG GAG TTC ATG GAT GAA GCT CTG ATC ATG GCA AGT ATG	2358
	Lys Ala Asn Val Glu Phe Met Asp Glu Ala Leu Ile Met Ala Ser Met	
	760 765 770 775	
10	GAT CAT CCA CAC CTA GTC CGG TTG CTG GGT GTG TGT CTG AGC CCA ACC	2406
	Asp His Pro His Leu Val Arg Leu Leu Gly Val Cys Leu Ser Pro Thr	
	780 785 790	
	ATC CAG CTG GTT ACT CAA CTT ATG CCC CAT GGC TGC CTG TTG GAG TAT	2454
	Ile Gln Leu Val Thr Gln Leu Met Pro His Gly Cys Leu Leu Glu Tyr	
	795 800 805	
15	GTC CAC GAG CAC AAG GAT AAC ATT GGA TCA CAA CTG CTG CTT AAC TGG	2502
	Val His Glu His Lys Asp Asn Ile Gly Ser Gln Leu Leu Leu Asn Trp	
	810 815 820	
20	TGT GTC CAG ATA GCT AAG GGA ATG ATG TAC CTG GAA GAA AGA CGA CTC	2550
	Cys Val Gln Ile Ala Lys Gly Met Met Tyr Leu Glu Glu Arg Arg Leu	
	825 830 835	
	GTT CAT CGG GAT TTG GCA GCC CGT AAT GTC TTA GTG AAA TCT CCA AAC	2598
	Val His Arg Asp Leu Ala Ala Arg Asn Val Leu Val Lys Ser Pro Asn	
	840 845 850 855	
25	CAT GTG AAA ATC ACA GAT TTT GGG CTA GCC AGA CTC TTG GAA GGA GAT	2646
	His Val Lys Ile Thr Asp Phe Gly Leu Ala Arg Leu Leu Glu Gly Asp	
	860 865 870	
30	GAA AAA GAG TAC AAT GCT GAT GGA GGA AAG ATG CCA ATT AAA TGG ATG	2694
	Glu Lys Glu Tyr Asn Ala Asp Gly Lys Lys Met Pro Ile Lys Trp Met	
	875 880 885	
	GCT CTG GAG TGT ATA CAT TAC AGG AAA TTC ACC CAT CAG AGT GAC GTT	2742
	Ala Leu Glu Cys Ile His Tyr Arg Lys Phe Thr His Gln Ser Asp Val	
	890 895 900	
35	TGG AGC TAT GGA GTT ACT ATA TGG GAA CTG ATG ACC TTT GGA GGA AAA	2790
	Trp Ser Tyr Gly Val Thr Ile Trp Glu Leu Met Thr Phe Gly Gly Lys	
	905 910 915	
40	CCC TAT GAT GGA ATT CCA ACG CGA GAA ATC CCT GAT TTA TTA GAG AAA	2838
	Pro Tyr Asp Gly Ile Pro Thr Arg Glu Ile Pro Asp Leu Leu Glu Lys	
	920 925 930 935	
	GGA GAA CGT TTG CCT CAG CCT CCC ATC TGC ACT ATT GAC GTT TAC ATG	2886
	Gly Glu Arg Leu Pro Gln Pro Pro Ile Cys Thr Ile Asp Val Tyr Met	
	940 945 950	
45	GTC ATG GTC AAA TGT TGG ATG ATT GAT GCT GAC AGT AGA CCT AAA TTT	2934
	Val Met Val Lys Cys Trp Met Ile Asp Ala Asp Ser Arg Pro Lys Phe	
	955 960 965	
50		
55		

	AAG GAA CTG GCT GCT GAG TTT TCA AGG ATG GCT CGA GAC CCT CAA AGA	2982
	Lys Glu Leu Ala Ala Glu Phe Ser Arg Met Ala Arg Asp Pro Gln Arg	
	970 975 980	
5	TAC CTA GTT ATT CAG GGT GAT GAT CGT ATG AAG CTT CCC AGT CCA AAT	3030
	Tyr Leu Val Ile Gln Gly Asp Asp Arg Met Lys Leu Pro Ser Pro Asn	
	985 990 995	
	GAC AGC AAG TTC TTT CAG AAT CTC TTG GAT GAA GAG GAT TTG GAA GAT	3078
	Asp Ser Lys Phe Phe Gln Asn Leu Leu Asp Glu Glu Asp Leu Glu Asp	
10	1000 1005 1010 1015	
	ATG ATG GAT GCT GAG GAG TAC TTG GTC CCT CAG GCT TTC AAC ATC CCA	3126
	Met Met Asp Ala Glu Glu Tyr Leu Val Pro Gln Ala Phe Asn Ile Pro	
	1020 1025 1030	
15	CCT CCC ATC TAT ACT TCC AGA GCA AGA ATT GAC TCG AAT AGG AGT GTA	3174
	Pro Pro Ile Tyr Thr Ser Arg Ala Arg Ile Asp Ser Asn Arg Ser Val	
	1035 1040 1045	
	AGA AAT AAT TAT ATA CAC ATA TCA TAT TCT TTC TGAGATATAA AATCATGTAA	3227
	Arg Asn Asn Tyr Ile His Ile Ser Tyr Ser Phe	
20	1050 1055	
	TAGTTCATAA GCACTAACAT TTCAAAATAA TTATATAGCT CAAATCAAAG TGATGCCTAG	3287
	ATTAAAAATA TACCATACCC ACAAAGATG TGCCAATCTT GCTATATGTA GTTAATTTTG	3347
	GAAGACAAGC ATGGACAATA CAACATGTAC TCTGAAATAC CTTCAAGATT TCAGAAGCAA	3407
25	AACATTTTCC TCATCTTAAT TTATTTAAAA CAAATCTTAA CTTTAAAAAA CAATTCCAAC	3467
	TAATAAAACC ATTATGTGTA TATAAATAAA TGAAAATTCC TACCAAGTAG GCTTTCTACT	3527
	TTTCTTTCTT AAAAAGATAT TATGATATAT TAGTCAAGAA GTAATACAAG TATAAATCTC	3587
30	TTTCACTTAT TTAAGAAAAA TTAAATATTT TCTGTCAAGT TGAAGTAGAA ACACAGAAAA	3647
	CCGTGCAGTC CTTTGAACCT AATCACATCG AAAAGGCTGC TGAGAAGTAG ATTTTGTGTT	3707
	TTAAGAAGTA GATTAAAGTT TTGAAGGAAG TTTCTGAAAA CACTTTACAT TTAAATGTT	3767
35	AAACCTACTC TATATGAATT CCATTCTTTC TTTGAAAGCT GTCAAATCCA TGCATTTATT	3827
	TTTATAAATT CATTCCTCAT ACATTCAACA TATATTGACT ACCACTGTAT GTGAAGCATT	3887
	AGTATACATT TAAGACTCAA AGAATTTTGA TACAACCTCT GCTTTCAAGA AGTGAAAACC	3947
40	TTAATCAAAG AATCATACAG ATAGAGGGAC TGCATAGTAA GTGCTGTAAT CCAGTATTCA	4007
	CTGACCAGTA CGGAGCATGA AGAAGTAGTA AATTTGTGTC TGTAATCAGT TTCTTCCATT	4067
	GATAAGATAT AAACATGATG CTTAATTTT TCTAGAAGAT AATTCTTTTC TCTTAATCTA	4127
45	AGAACATTAT CATAGCTAGT AGAACCGACA GCATCCGATT TCTCTTGACC ATAGCCATAA	4187
	GAATATCTTC AACTTGCTGC TCATTATCTA ACAAACATAA TTTTCTTTAT TTCATATTGA	4247

50

55

TTGTAATAAG TAATATCCCC CTGGAAGTTT ACTATTCAAC ACATATATGT TAACCTCCTT 4307  
 AATTCCTTAA ACAAACCTCA TGAGGTTCTA TTATTATCAT CCCCTTCTTT CAAAGGAAGA 4367  
 5 AACTTGCCAC AGAGAAGTCA GGTGATATGA CTGGTGTAC ACAGCTAGTC AGTGAAGAG 4427  
 AGGAATAAGT AATCTAGATA TCTGCCTACT ACACGTAGG TTTGCTTCAA AGTTACTGAA 4487  
 GYCATGTTAT TTCCATGATG TGATTAGAGT CTGGGACTTG TCTTGTTTGG GAAATTTCCC 4547  
 10 AGGTGGTTTT CTTATAAAAT GCATCTCAAA TCTGCTCTAC ACCTTTTACT CATCTACCTC 4607  
 CATTAGAAG ATCTGATATG GAAAGAGACA AAGATGGAGA CCTCAATTAT TTTTCTTTT 4667  
 CTGTTAAAAA TATTATAGTA CAACTGAAAC TTATCACATG CCAATGGGGA ATAGATAACT 4727  
 15 AAAAGTTTAA AATTAGATCA ATGGATAGGT AAATGAATAA TCNTTCTTTT GCTGTGAGA 4787  
 GGGGAAGGAA AAGCGGTAA GGTGGTATAA AGGAGGCTCC TCTGTACACT TGCAAAATGA 4847  
 TCAAATTATA TACCCTTGTA TTTATAATTT TAAGTGACAA ATTCACTACT TCTGGTTACA 4907  
 20 ACAGTGAAAT TTAAAAAAA ATAGTTTTTC TTTCTTAGCT TGCAATGCTA TAAATCTTTT 4967  
 TCTTTTATA AGAATTCTTA CATTTCAGCT TTTTGTTCAT TTTAATTTAT AATTCTCAGT 5027  
 GCAAGAAATT CTTAATAAAG GTTTGAGCTA GCTAGATGGA ATTATTGAGA CAAAGTCTAA 5087  
 25 ATCACCCGTG GACTTATTTG ACCTTTAGCC ATCATTCTT ATTCCACATT ATAAAAAAT 5147  
 GTTACCTGTA GATTTCCTTT TACTTTTTCA GTCCTTGGAA AAGAAATGGT GATTAAATAT 5207  
 CATTATATCA TTTTATGTTT AGGCATTTAA AAAGCTTTAT TTGTCATCTA TATTGCTCTA 5267  
 30 ATAGTTTTCA GTCTGGCTTT ACGTAACTTT TACGGAAATT TCTAACATGT ACAAATGCCA 5327  
 TGTTCTCCTT TTCTTTCCTA CATGGCTGAA TTAGAAAACA AATTACTTCC ATTTTAAGTT 5387  
 TGGCTAAAT AGAAAACAAA TTAATACCAT TTTAAGTTT GTGGCTAAAT AACGTGCTAA 5447  
 35 GGGAACATCT TAAAAAGTGA ATTTTGATCA AATATTTCTT AAGCATATGT GATAGACTTT 5507  
 GAAACCAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA 5555

## (2) INFORMATION FOR SEQ ID NO:4:

40 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 1058 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Lys Pro Ala Thr Gly Leu Trp Val Trp Val Ser Leu Leu Val Ala  
 1 5 10 15

50

55

Ala Gly Thr Val Gln Pro Ser Asp Ser Gln Ser Val Cys Ala Gly Thr  
                   20                  25                  30  
 5 Glu Asn Lys Leu Ser Ser Leu Ser Asp Leu Glu Gln Gln Tyr Arg Ala  
           35                  40                  45  
 Leu Arg Lys Tyr Tyr Glu Asn Cys Glu Val Val Met Gly Asn Leu Glu  
           50                  55                  60  
 10 Ile Thr Ser Ile Glu His Asn Arg Asp Leu Ser Phe Leu Arg Ser Val  
       65                  70                  75                  80  
 Arg Glu Val Thr Gly Tyr Val Leu Val Ala Leu Asn Gln Phe Arg Tyr  
           85                  90                  95  
 15 Leu Pro Leu Glu Asn Leu Arg Ile Ile Arg Gly Thr Lys Leu Tyr Glu  
           100                  105                  110  
 Asp Arg Tyr Ala Leu Ala Ile Phe Leu Asn Tyr Arg Lys Asp Gly Asn  
           115                  120                  125  
 20 Phe Gly Leu Gln Glu Leu Gly Leu Lys Asn Leu Thr Glu Ile Leu Asn  
       130                  135                  140  
 Gly Gly Val Tyr Val Asp Gln Asn Lys Phe Leu Cys Tyr Ala Asp Thr  
       145                  150                  155                  160  
 25 Ile His Trp Gln Asp Ile Val Arg Asn Pro Trp Pro Ser Asn Leu Thr  
           165                  170                  175  
 Leu Val Ser Thr Asn Gly Ser Ser Gly Cys Gly Arg Cys His Lys Ser  
           180                  185                  190  
 30 Cys Thr Gly Arg Cys Trp Gly Pro Thr Glu Asn His Cys Gln Thr Leu  
       195                  200                  205  
 Thr Arg Thr Val Cys Ala Glu Gln Cys Asp Gly Arg Cys Tyr Gly Pro  
       210                  215                  220  
 35 Tyr Val Ser Asp Cys Cys His Arg Glu Cys Ala Gly Gly Cys Ser Gly  
       225                  230                  235                  240  
 Pro Lys Asp Thr Asp Cys Phe Ala Cys Met Asn Phe Asn Asp Ser Gly  
           245                  250                  255  
 40 Ala Cys Val Thr Gln Cys Pro Gln Thr Phe Val Tyr Asn Pro Thr Thr  
           260                  265                  270  
 45 Phe Gln Leu Glu His Asn Phe Asn Ala Lys Tyr Thr Tyr Gly Ala Phe  
           275                  280                  285  
 Cys Val Lys Lys Cys Pro His Asn Phe Val Val Asp Ser Ser Ser Cys  
       290                  295                  300  
 50 Val Arg Ala Cys Pro Ser Ser Lys Met Glu Val Glu Glu Asn Gly Ile  
       305                  310                  315                  320  
 55



Cys Asn Gly Pro Thr Ser His Asp Cys Ile Tyr Tyr Pro Trp Thr Gly  
 625 630 635 640  
 5 His Ser Thr Leu Pro Gln His Ala Arg Thr Pro Leu Ile Ala Ala Gly  
 645 650 655  
 Val Ile Gly Gly Leu Phe Ile Leu Val Ile Val Gly Leu Thr Phe Ala  
 660 665 670  
 10 Val Tyr Val Arg Arg Lys Ser Ile Lys Lys Lys Arg Ala Leu Arg Arg  
 675 680 685  
 Phe Leu Glu Thr Glu Leu Val Glu Pro Leu Thr Pro Ser Gly Thr Ala  
 690 695 700  
 15 Pro Asn Gln Ala Gln Leu Arg Ile Leu Lys Glu Thr Glu Leu Lys Arg  
 705 710 715 720  
 Val Lys Val Leu Gly Ser Gly Ala Phe Gly Thr Val Tyr Lys Gly Ile  
 725 730 735  
 20 Trp Val Pro Glu Gly Glu Thr Val Lys Ile Pro Val Ala Ile Lys Ile  
 740 745 750  
 Leu Asn Glu Thr Thr Gly Pro Lys Ala Asn Val Glu Phe Met Asp Glu  
 755 760 765  
 25 Ala Leu Ile Met Ala Ser Met Asp His Pro His Leu Val Arg Leu Leu  
 770 775 780  
 Gly Val Cys Leu Ser Pro Thr Ile Gln Leu Val Thr Gln Leu Met Pro  
 785 790 795 800  
 30 His Gly Cys Leu Leu Glu Tyr Val His Glu His Lys Asp Asn Ile Gly  
 805 810 815  
 Ser Gln Leu Leu Leu Asn Trp Cys Val Gln Ile Ala Lys Gly Met Met  
 820 825 830  
 35 Tyr Leu Glu Glu Arg Arg Leu Val His Arg Asp Leu Ala Ala Arg Asn  
 835 840 845  
 Val Leu Val Lys Ser Pro Asn His Val Lys Ile Thr Asp Phe Gly Leu  
 850 855 860  
 Ala Arg Leu Leu Glu Gly Asp Glu Lys Glu Tyr Asn Ala Asp Gly Gly  
 865 870 875 880  
 45 Lys Met Pro Ile Lys Trp Met Ala Leu Glu Cys Ile His Tyr Arg Lys  
 885 890 895  
 Phe Thr His Gln Ser Asp Val Trp Ser Tyr Gly Val Thr Ile Trp Glu  
 900 905 910  
 50 Leu Met Thr Phe Gly Gly Lys Pro Tyr Asp Gly Ile Pro Thr Arg Glu  
 915 920 925

55

Ile Pro Asp Leu Leu Glu Lys Gly Glu Arg Leu Pro Gln Pro Pro Ile  
 930 935 940  
 Cys Thr Ile Asp Val Tyr Met Val Met Val Lys Cys Trp Met Ile Asp  
 8 945 950 955 960  
 Ala Asp Ser Arg Pro Lys Phe Lys Glu Leu Ala Ala Glu Phe Ser Arg  
 965 970 975  
 Met Ala Arg Asp Pro Gln Arg Tyr Leu Val Ile Gln Gly Asp Asp Arg  
 10 980 985 990  
 Met Lys Leu Pro Ser Pro Asn Asp Ser Lys Phe Phe Gln Asn Leu Leu  
 995 1000 1005  
 Asp Glu Glu Asp Leu Glu Asp Met Met Asp Ala Glu Glu Tyr Leu Val  
 15 1010 1015 1020  
 Pro Gln Ala Phe Asn Ile Pro Pro Pro Ile Tyr Thr Ser Arg Ala Arg  
 1025 1030 1035 1040  
 Ile Asp Ser Asn Arg Ser Val Arg Asn Asn Tyr Ile His Ile Ser Tyr  
 20 1045 1050 1055  
 Ser Phe

## (2) INFORMATION FOR SEQ ID NO:5:

25 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 3321 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: unknown

30 (ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 156..1782

35

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

CATTAGCTGC AATTGATCAA GTGACTGAGA GAAGGGCAAC ATTCCATGCA ACAGTATAGT 60  
 40 GGTATGGAAA GCCCTGGATG TTGAAATCTA GCTTCAAAAA GCCTGTCTGG AAATGTAGTT 120  
 AATTGGATGA AGTGAGAAGA GATAAAACCA GAGAG GAA GCT CTG ATC ATG GCA 173  
 Glu Ala Leu Ile Met Ala  
 1 5  
 45 AGT ATG GAT CAT CCA CAC CTA GTC CGG TTG CTG GGT GTG TGT CTG AGC 221  
 Ser Met Asp His Pro His Leu Val Arg Leu Leu Gly Val Cys Leu Ser  
 10 15 20

50

55

	CCA ACC ATC CAG CTG GTT ACT CAA CTT ATG CCC CAT GGC TGC CTG TTG	269
	Pro Thr Ile Gln Leu Val Thr Gln Leu Met Pro His Gly Cys Leu Leu	
	25 30 35	
5	GAG TAT GTC CAC GAG CAC AAG GAT AAC ATT GGA TCA CAA CTG CTG CTT	317
	Glu Tyr Val His Glu His Lys Asp Asn Ile Gly Ser Gln Leu Leu Leu	
	40 45 50	
10	AAC TGG TGT GTC CAG ATA GCT AAG GGA ATG ATG TAC CTG GAA GAA AGA	365
	Asn Trp Cys Val Gln Ile Ala Lys Gly Met Met Tyr Leu Glu Glu Arg	
	55 60 65 70	
	CGA CTC GTT CAT CGG GAT TTG GCA GCC CGT AAT GTC TTA GTG AAA TCT	413
	Arg Leu Val His Arg Asp Leu Ala Ala Arg Asn Val Leu Val Lys Ser	
	75 80 85	
15	CCA AAC CAT GTG AAA ATC ACA GAT TTT GGG CTA GCC AGA CTC TTG GAA	461
	Pro Asn His Val Lys Ile Thr Asp Phe Gly Leu Ala Arg Leu Leu Glu	
	90 95 100	
20	GGA GAT GAA AAA GAG TAC AAT GCT GAT GGA GGA AAG ATG CCA ATT AAA	509
	Gly Asp Glu Lys Glu Tyr Asn Ala Asp Gly Gly Lys Met Pro Ile Lys	
	105 110 115	
	TGG ATG GCT CTG GAG TGT ATA CAT TAC AGG AAA TTC ACC CAT CAG AGT	557
	Trp Met Ala Leu Glu Cys Ile His Tyr Arg Lys Phe Thr His Gln Ser	
	120 125 130	
25	GAC GTT TGG AGC TAT GGA GTT ACT ATA TGG GAA CTG ATG ACC TTT GGA	605
	Asp Val Trp Ser Tyr Gly Val Thr Ile Trp Glu Leu Met Thr Phe Gly	
	135 140 145 150	
30	GGA AAA CCC TAT GAT GGA ATT CCA ACG CGA GAA ATC CCT GAT TTA TTA	653
	Gly Lys Pro Tyr Asp Gly Ile Pro Thr Arg Glu Ile Pro Asp Leu Leu	
	155 160 165	
	GAG AAA GGA GAA CGT TTG CCT CAG CCT CCC ATC TGC ACT ATT GAC GTT	701
	Glu Lys Gly Glu Arg Leu Pro Gln Pro Pro Ile Cys Thr Ile Asp Val	
	170 175 180	
35	TAC ATG GTC ATG GTC AAA TGT TGG ATG ATT GAT GCT GAC AGT AGA CCT	749
	Tyr Met Val Met Val Lys Cys Trp Met Ile Asp Ala Asp Ser Arg Pro	
	185 190 195	
40	AAA TTT AAG GAA CTG GCT GCT GAG TTT TCA AGG ATG GCT CGA GAC CCT	797
	Lys Phe Lys Glu Leu Ala Ala Glu Phe Ser Arg Met Ala Arg Asp Pro	
	200 205 210	
	CAA AGA TAC CTA GTT ATT CAG GGT GAT GAT CGT ATG AAG CTT CCC AGT	845
	Gln Arg Tyr Leu Val Ile Gln Gly Asp Asp Arg Met Lys Leu Pro Ser	
	215 220 225 230	
45	CCA AAT GAC AGC AAG TTC TTT CAG AAT CTC TTG GAT GAA GAG GAT TTG	893
	Pro Asn Asp Ser Lys Phe Phe Gln Asn Leu Leu Asp Glu Glu Asp Leu	
	235 240 245	

50

55



	GAA	GAT	ATG	ATG	GAT	GCT	GAG	GAG	TAC	TTG	GTC	CCT	CAG	GCT	TTC	AAC	941
	Glu	Asp	Met	Met	Asp	Ala	Glu	Glu	Tyr	Leu	Val	Pro	Gln	Ala	Phe	Asn	
				250					255								
5	ATC	CCA	CCT	CCC	ATC	TAT	ACT	TCC	AGA	GCA	AGA	ATT	GAC	TCG	AAT	AGG	989
	Ile	Pro	Pro	Pro	Ile	Tyr	Thr	Ser	Arg	Ala	Arg	Ile	Asp	Ser	Asn	Arg	
			265					270					275				
10	AGT	GAA	ATT	GGA	CAC	AGC	CCT	CCT	CCT	GCC	TAC	ACC	CCC	ATG	TCA	GGA	1037
	Ser	Glu	Ile	Gly	His	Ser	Pro	Pro	Pro	Ala	Tyr	Thr	Pro	Met	Ser	Gly	
			280				285					290					
	AAC	CAG	TTT	GTA	TAC	CGA	GAT	GGA	GGT	TTT	GCT	GCT	GAA	CAA	GGA	GTG	1085
	Asn	Gln	Phe	Val	Tyr	Arg	Asp	Gly	Gly	Phe	Ala	Ala	Glu	Gln	Gly	Val	
						300					305					310	
15	TCT	GTG	CCC	TAC	AGA	GCC	CCA	ACT	AGC	ACA	ATT	CCA	GAA	GCT	CCT	GTG	1133
	Ser	Val	Pro	Tyr	Arg	Ala	Pro	Thr	Ser	Thr	Ile	Pro	Glu	Ala	Pro	Val	
					315					320					325		
20	GCA	CAG	GGT	GCT	ACT	GCT	GAG	ATT	TTT	GAT	GAC	TCC	TGC	TGT	AAT	GGC	1181
	Ala	Gln	Gly	Ala	Thr	Ala	Glu	Ile	Phe	Asp	Asp	Ser	Cys	Cys	Asn	Gly	
				330					335					340			
	ACC	CTA	CGC	AAG	CCA	GTG	GCA	CCC	CAT	GTC	CAA	GAG	GAC	AGT	AGC	ACC	1229
	Thr	Leu	Arg	Lys	Pro	Val	Ala	Pro	His	Val	Gln	Glu	Asp	Ser	Ser	Thr	
				345				350					355				
25	CAG	AGG	TAC	AGT	GCT	GAC	CCC	ACC	GTG	TTT	GCC	CCA	GAA	CGG	AGC	CCA	1277
	Gln	Arg	Tyr	Ser	Ala	Asp	Pro	Thr	Val	Phe	Ala	Pro	Glu	Arg	Ser	Pro	
							365					370					
30	CGA	GGA	GAG	CTG	GAT	GAG	GAA	GGT	TAC	ATG	ACT	CCT	ATG	CGA	GAC	AAA	1325
	Arg	Gly	Glu	Leu	Asp	Glu	Glu	Gly	Tyr	Met	Thr	Pro	Met	Arg	Asp	Lys	
						380					385					390	
	CCC	AAA	CAA	GAA	TAC	CTG	AAT	CCA	GTG	GAG	GAG	AAC	CCT	TTT	GTT	TCT	1373
	Pro	Lys	Gln	Glu	Tyr	Leu	Asn	Pro	Val	Glu	Glu	Asn	Pro	Phe	Val	Ser	
					395					400					405		
35	CGG	AGA	AAA	AAT	GGA	GAC	CTT	CAA	GCA	TTG	GAT	AAT	CCC	GAA	TAT	CAC	1421
	Arg	Arg	Lys	Asn	Gly	Asp	Leu	Gln	Ala	Leu	Asp	Asn	Pro	Glu	Tyr	His	
				410					415					420			
	AAT	GCA	TCC	AAT	GGT	CCA	CCC	AAG	GCC	GAG	GAT	GAG	TAT	GTG	AAT	GAG	1469
	Asn	Ala	Ser	Asn	Gly	Pro	Pro	Lys	Ala	Glu	Asp	Glu	Tyr	Val	Asn	Glu	
				425				430					435				
40	CCA	CTG	TAC	CTC	AAC	ACC	TTT	GCC	AAC	ACC	TTG	GGA	AAA	GCT	GAG	TAC	1517
	Pro	Leu	Tyr	Leu	Asn	Thr	Phe	Ala	Asn	Thr	Leu	Gly	Lys	Ala	Glu	Tyr	
						445						450					
45	CTG	AAG	AAC	AAC	ATA	CTG	TCA	ATG	CCA	GAG	AAG	GCC	AAG	AAA	GCG	TTT	1565
	Leu	Lys	Asn	Asn	Ile	Leu	Ser	Met	Pro	Glu	Lys	Ala	Lys	Lys	Ala	Phe	
						460					465					470	
50																	
55																	

	GAC AAC CCT GAC TAC TGG AAC CAC AGC CTG CCA CCT CGG AGC ACC CTT Asp Asn Pro Asp Tyr Trp Asn His Ser Leu Pro Pro Arg Ser Thr Leu 475 480 485	1613
5	CAG CAC CCA GAC TAC CTG CAG GAG TAC AGC ACA AAA TAT TTT TAT AAA Gln His Pro Asp Tyr Leu Gln Glu Tyr Ser Thr Lys Tyr Phe Tyr Lys 490 495 500	1661
	CAG AAT GGG CGG ATC CGG CCT ATT GTG GCA GAG AAT CCT GAA TAC CTC Gln Asn Gly Arg Ile Arg Pro Ile Val Ala Glu Asn Pro Glu Tyr Leu 505 510 515	1709
10	TCT GAG TTC TCC CTG AAG CCA GGC ACT GTG CTG CCG CCT CCA CCT TAC Ser Glu Phe Ser Leu Lys Pro Gly Thr Val Leu Pro Pro Pro Pro Tyr 520 525 530	1757
15	AGA CAC CGG AAT ACT GTG GTG TAAGCTCAGT TGTGCTTTT TAGGTGGAGA Arg His Arg Asn Thr Val Val 535 540	1808
	GACACACCTG CTCCAATTTC CCCACCCCCC TCTCTTTCTC TGGTGGTCTT CCTTCTACCC	1868
20	CAAGGCCAGT AGTTTTCACA CTTCCCACTG GAAGATACAG AGATGCAATG ATAGTTATGT GCTTACCTAA CTTGAAACATT AGAGGGAAG ACTGAAAGAG AAAGATAGGA GGAACCACAA TGTTTCTTCA TTTCTCTGCA TGGGTGGTC AGGAGAATGA AACAGCTAGA GAAGGACCAG	1928 1988 2048
25	AAAATGTAAG GCAATGCTGC CTAATATCAA ACTAGCTGTC ACTTTTTTTC TTTTCTTTT TCTTTCTTTG TTTCTTTCTT CCTCTCTTT TTTTTTTTTT TTTTAAAGCA GATGGTTGAA ACACCCATGC TATCTGTTCC TATCTGCAGG AACTGATGTG TGCATATTTA GCATCCCTGG	2108 2168 2228
30	AAATCATAAT AAAGTTTCCA TTAGAACAAA AGAATAACAT TTTCTATAAC ATATGATAGT GTCTGAAATT GAGAATCCAG TTTCTTTCCC CAGCAGTTTC TGTCTAGCA AGTAAGAATG GCCAACTCAA CTTTCATAAT TTAAAAATCT CCATTAAAGT TATAACTAGT AATTATGTTT	2288 2348 2408
35	TCAACACTTT TTGGTTTTTT TCATTTTGTT TTGCTCTGAC CGATTCCCTT ATATTTGCTC CCCTATTTTT GGCTTTAATT TCTAATTGCA AAGATGTTTA CATCAAAGCT TCTTCACAGA ATTTAAGCAA GAAATATTTT AATATAGTGA AATGGCCACT ACTTTAAGTA TACAATCTTT	2468 2528 2588
40	AAAATAAGAA AGGGAGGCTA ATATTTTTC TGTATCAAA TTATCTTCAC CCTCATCCTT TACATTTTTT AACATTTTTT TTTCTCCATA AATGACACTA CTTGATAGGC CGTTGGTTGT CTGAAGAGTA GAAGGGAAC TAAGAGACAG TTCTCTGTGG TTCAGGAAAA CTAATGATAC	2648 2708 2768
45	TTTCAGGGGT GGCCCAATGA GGGAAATCAT TGAAGTGGAA GAAACACACT GGATTGGGTA TGTCTACCTG GCAGATACTC AGAAATGTAG TTGCACTTA AGCTGTAATT TTATTTGTTT	2828 2888

50

55

TTTTCTGAA CTCCATTTTG GATTTTGAAT CAAGCAATAT GGAAGCAACC AGCAAATTAA 2948  
 CTAATTAAAG TACATTTTAA AAAAAAGAGC TAAGATAAAG ACTGTGGAAA TGCCAAACCA 3008  
 6 AGCAAATTAG GAACCTTGCA ACGGTATCCA GGGACTATGA TGAGAGGCCA GCACATTATC 3068  
 TTCATATGTC ACCTTTGCTA CGCAAGGAAA TTTGTTTCACT TCGTATACTT CGTAAGAAGG 3128  
 AATGCGAGTA AGGATTGGCT TGAATTCAT GGAATTTCTA GTATGAGACT ATTTATATGA 3188  
 10 AGTAGAAGGT AACTCTTTGC ACATAAATTG GTATAATAAA AAGAAAAACA CAAACATTCA 3248  
 AAGCTTAGGG ATAGGTCCTT GGGTCAAAAG TTGTAAATAA ATGTGAAACA TCTTCTCAAA 3308  
 AAAAAAAAAA AAA 3321

15 (2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 541 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

25 Glu Ala Leu Ile Met Ala Ser Met Asp His Pro His Leu Val Arg Leu  
 1 5 10 15  
 Leu Gly Val Cys Leu Ser Pro Thr Ile Gln Leu Val Thr Gln Leu Met  
 20 25 30  
 30 Pro His Gly Cys Leu Leu Glu Tyr Val His Glu His Lys Asp Asn Ile  
 35 40 45  
 Gly Ser Gln Leu Leu Leu Asn Trp Cys Val Gln Ile Ala Lys Gly Met  
 50 55 60  
 35 Met Tyr Leu Glu Glu Arg Arg Leu Val His Arg Asp Leu Ala Ala Arg  
 65 70 75 80  
 Asn Val Leu Val Lys Ser Pro Asn His Val Lys Ile Thr Asp Phe Gly  
 85 90 95  
 40 Leu Ala Arg Leu Leu Glu Gly Asp Glu Lys Glu Tyr Asn Ala Asp Gly  
 100 105 110  
 Gly Lys Met Pro Ile Lys Trp Met Ala Leu Glu Cys Ile His Tyr Arg  
 115 120 125  
 45 Lys Phe Thr His Gln Ser Asp Val Trp Ser Tyr Gly Val Thr Ile Trp  
 130 135 140  
 Glu Leu Met Thr Phe Gly Gly Lys Pro Tyr Asp Gly Ile Pro Thr Arg  
 145 150 155 160

50

55

Glu Ile Pro Asp Leu Leu Glu Lys Gly Glu Arg Leu Pro Gln Pr Pr  
 165 170 175  
 5 Ile Cys Thr Ile Asp Val Tyr Met Val Met Val Lys Cys Trp Met Ile  
 180 185 190  
 Asp Ala Asp Ser Arg Pro Lys Phe Lys Glu Leu Ala Ala Glu Phe Ser  
 195 200 205  
 10 Arg Met Ala Arg Asp Pro Gln Arg Tyr Leu Val Ile Gln Gly Asp Asp  
 210 215 220  
 Arg Met Lys Leu Pro Ser Pro Asn Asp Ser Lys Phe Phe Gln Asn Leu  
 225 230 235 240  
 15 Leu Asp Glu Glu Asp Leu Glu Asp Met Met Asp Ala Glu Glu Tyr Leu  
 245 250 255  
 Val Pro Gln Ala Phe Asn Ile Pro Pro Pro Ile Tyr Thr Ser Arg Ala  
 260 265 270  
 20 Arg Ile Asp Ser Asn Arg Ser Glu Ile Gly His Ser Pro Pro Pro Ala  
 275 280 285  
 Tyr Thr Pro Met Ser Gly Asn Gln Phe Val Tyr Arg Asp Gly Gly Phe  
 290 295 300  
 25 Ala Ala Glu Gln Gly Val Ser Val Pro Tyr Arg Ala Pro Thr Ser Thr  
 305 310 315 320  
 Ile Pro Glu Ala Pro Val Ala Gln Gly Ala Thr Ala Glu Ile Phe Asp  
 325 330 335  
 30 Asp Ser Cys Cys Asn Gly Thr Leu Arg Lys Pro Val Ala Pro His Val  
 340 345 350  
 Gln Glu Asp Ser Ser Thr Gln Arg Tyr Ser Ala Asp Pro Thr Val Phe  
 355 360 365  
 35 Ala Pro Glu Arg Ser Pro Arg Gly Glu Leu Asp Glu Glu Gly Tyr Met  
 370 375 380  
 40 Thr Pro Met Arg Asp Lys Pro Lys Gln Glu Tyr Leu Asn Pro Val Glu  
 385 390 395 400  
 Glu Asn Pro Phe Val Ser Arg Arg Lys Asn Gly Asp Leu Gln Ala Leu  
 405 410 415  
 45 Asp Asn Pro Glu Tyr His Asn Ala Ser Asn Gly Pro Pro Lys Ala Glu  
 420 425 430  
 Asp Glu Tyr Val Asn Glu Pro Leu Tyr Leu Asn Thr Phe Ala Asn Thr  
 435 440 445  
 50 Leu Gly Lys Ala Glu Tyr Leu Lys Asn Asn Ile Leu Ser Met Pro Glu  
 450 455 460

55

Lys Ala Lys Lys Ala Phe Asp Asn Pr Asp Tyr Trp Asn His Ser Leu  
 465 470 475 480  
 5 Pro Pro Arg Ser Thr Leu Gln His Pro Asp Tyr Leu Gln Glu Tyr Ser  
 485 490 495  
 Thr Lys Tyr Phe Tyr Lys Gln Asn Gly Arg Ile Arg Pro Ile Val Ala  
 500 505 510  
 10 Glu Asn Pro Glu Tyr Leu Ser Glu Phe Ser Leu Lys Pro Gly Thr Val  
 515 520 525  
 Leu Pro Pro Pro Pro Tyr Arg His Arg Asn Thr Val Val  
 530 535 540  
 15 (2) INFORMATION FOR SEQ ID NO:7:  
 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 1210 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: unknown  
 (D) TOPOLOGY: unknown  
 20 (ii) MOLECULE TYPE: protein  
 25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:  
 Met Arg Pro Ser Gly Thr Ala Gly Ala Ala Leu Leu Ala Leu Leu Ala  
 1 5 10 15  
 30 Ala Leu Cys Pro Ala Ser Arg Ala Leu Glu Glu Lys Lys Val Cys Gln  
 20 25 30  
 Gly Thr Ser Asn Lys Leu Thr Gln Leu Gly Thr Phe Glu Asp His Phe  
 35 40 45  
 Leu Ser Leu Gln Arg Met Phe Asn Asn Cys Glu Val Val Leu Gly Asn  
 50 55 60  
 Leu Glu Ile Thr Tyr Val Gln Arg Asn Tyr Asp Leu Ser Phe Leu Lys  
 65 70 75 80  
 40 Thr Ile Gln Glu Val Ala Gly Tyr Val Leu Ile Ala Leu Asn Thr Val  
 85 90 95  
 Glu Arg Ile Pro Leu Glu Asn Leu Gln Ile Ile Arg Gly Asn Met Tyr  
 100 105 110  
 45 Tyr Glu Asn Ser Tyr Ala Leu Ala Val Leu Ser Asn Tyr Asp Ala Asn  
 115 120 125  
 Lys Thr Gly Leu Lys Glu Leu Pro Met Arg Asn Leu Gln Glu Ile Leu  
 130 135 140  
 50  
 55

His Gly Ala Val Arg Phe Ser Asn Asn Pro Ala Leu Cys Asn Val Glu  
 145 150 155 160  
 5 Ser Ile Gln Trp Arg Asp Ile Val Ser Ser Asp Phe Leu Ser Asn Met  
 165 170 175  
 Ser Met Asp Phe Gln Asn His Leu Gly Ser Cys Gln Lys Cys Asp Pro  
 180 185 190  
 10 Ser Cys Pro Asn Gly Ser Cys Trp Gly Ala Gly Glu Glu Asn Cys Gln  
 195 200 205  
 Lys Leu Thr Lys Ile Ile Cys Ala Gln Gln Cys Ser Gly Arg Cys Arg  
 210 215 220  
 15 Gly Lys Ser Pro Ser Asp Cys Cys His Asn Gln Cys Ala Ala Gly Cys  
 225 230 235 240  
 Thr Gly Pro Arg Glu Ser Asp Cys Leu Val Cys Arg Lys Phe Arg Asp  
 245 250 255  
 20 Glu Ala Thr Cys Lys Asp Thr Cys Pro Pro Leu Met Leu Tyr Asn Pro  
 260 265 270  
 Thr Thr Tyr Gln Met Asp Val Asn Pro Glu Gly Lys Tyr Ser Phe Gly  
 275 280 285  
 25 Ala Thr Cys Val Lys Lys Cys Pro Arg Asn Tyr Val Val Thr Asp His  
 290 295 300  
 Gly Ser Cys Val Arg Ala Cys Gly Ala Asp Ser Tyr Glu Met Glu Glu  
 305 310 315 320  
 30 Asp Gly Val Arg Lys Cys Lys Lys Cys Glu Gly Pro Cys Arg Lys Val  
 325 330 335  
 Cys Asn Gly Ile Gly Ile Gly Glu Phe Lys Asp Ser Leu Ser Ile Asn  
 340 345 350  
 35 Ala Thr Asn Ile Lys His Phe Lys Asn Cys Thr Ser Ile Ser Gly Asp  
 355 360 365  
 Leu His Ile Leu Pro Val Ala Phe Arg Gly Asp Ser Phe Thr His Thr  
 370 375 380  
 40 Pro Pro Leu Asp Pro Gln Glu Leu Asp Ile Leu Lys Thr Val Lys Glu  
 385 390 395 400  
 Ile Thr Gly Phe Leu Leu Ile Gln Ala Trp Pro Glu Asn Arg Thr Asp  
 405 410 415  
 45 Leu His Ala Phe Glu Asn Leu Glu Ile Ile Arg Gly Arg Thr Lys Gln  
 420 425 430  
 His Gly Gln Phe Ser Leu Ala Val Val Ser Leu Asn Ile Thr Ser Leu  
 435 440 445  
 50  
 55

Gly Leu Arg Ser Leu Lys Glu Ile Ser Asp Gly Asp Val Ile Ile Ser  
 450 455 460  
 6 Gly Asn Lys Asn Leu Cys Tyr Ala Asn Thr Ile Asn Trp Lys Lys Leu  
 465 470 475 480  
 Phe Gly Thr Ser Gly Gln Lys Thr Lys Ile Ile Ser Asn Arg Gly Glu  
 485 490 495  
 10 Asn Ser Cys Lys Ala Thr Gly Gln Val Cys His Ala Leu Cys Ser Pro  
 500 505 510  
 Glu Gly Cys Trp Gly Pro Glu Pro Arg Asp Cys Val Ser Cys Arg Asn  
 515 520 525  
 15 Val Ser Arg Gly Arg Glu Cys Val Asp Lys Cys Lys Leu Leu Glu Gly  
 530 535 540  
 Glu Pro Arg Glu Phe Val Glu Asn Ser Glu Cys Ile Gln Cys His Pro  
 545 550 555 560  
 20 Glu Cys Leu Pro Gln Ala Met Asn Ile Thr Cys Thr Gly Arg Gly Pro  
 565 570 575  
 Asp Asn Cys Ile Gln Cys Ala His Tyr Ile Asp Gly Pro His Cys Val  
 580 585 590  
 25 Lys Thr Cys Pro Ala Gly Val Met Gly Glu Asn Asn Thr Leu Val Trp  
 595 600 605  
 Lys Tyr Ala Asp Ala Gly His Val Cys His Leu Cys His Pro Asn Cys  
 610 615 620  
 30 Thr Tyr Gly Cys Thr Gly Pro Gly Leu Glu Gly Cys Pro Thr Asn Gly  
 625 630 635 640  
 Pro Lys Ile Pro Ser Ile Ala Thr Gly Met Val Gly Ala Leu Leu Leu  
 645 650 655  
 35 Leu Leu Val Val Ala Leu Gly Ile Gly Leu Phe Met Arg Arg Arg His  
 660 665 670  
 Ile Val Arg Lys Arg Thr Leu Arg Arg Leu Leu Gln Glu Arg Glu Leu  
 675 680 685  
 40 Val Glu Pro Leu Thr Pro Ser Gly Glu Ala Pro Asn Gln Ala Leu Leu  
 690 695 700  
 Arg Ile Leu Lys Glu Thr Glu Phe Lys Lys Ile Lys Val Leu Gly Ser  
 705 710 715 720  
 Gly Ala Phe Gly Thr Val Tyr Lys Gly Leu Trp Ile Pro Glu Gly Glu  
 725 730 735  
 50 Lys Val Lys Ile Pro Val Ala Ile Lys Glu Leu Arg Glu Ala Thr Ser  
 740 745 750

55

Pro Lys Ala Asn Lys Glu Ile Leu Asp Glu Ala Tyr Val Met Ala Ser  
 755 760 765  
 5 Val Asp Asn Pr His Val Cys Arg Leu Leu Gly Ile Cys Leu Thr Ser  
 770 775 780  
 Thr Val Gln Leu Ile Thr Gln Leu Met Pro Phe Gly Cys Leu Leu Asp  
 785 790 795 800  
 10 Tyr Val Arg Glu His Lys Asp Asn Ile Gly Ser Gln Tyr Leu Leu Asn  
 805 810 815  
 Trp Cys Val Gln Ile Ala Lys Gly Met Met Tyr Leu Glu Asp Arg Arg  
 820 825 830  
 15 Leu Val His Arg Asp Leu Ala Ala Arg Asn Val Leu Val Lys Thr Pro  
 835 840 845  
 Gln His Val Lys Ile Thr Asp Phe Gly Leu Ala Lys Leu Leu Gly Ala  
 850 855 860  
 20 Glu Glu Lys Glu Tyr His Ala Glu Gly Gly Lys Val Pro Ile Lys Trp  
 865 870 875 880  
 Met Ala Leu Glu Ser Ile Leu His Arg Ile Tyr Thr His Gln Ser Asp  
 885 890 895  
 25 Val Trp Ser Tyr Gly Val Thr Val Trp Glu Leu Met Thr Phe Gly Ser  
 900 905 910  
 Lys Pro Tyr Asp Gly Ile Pro Ala Ser Glu Ile Ser Ser Ile Leu Glu  
 915 920 925  
 30 Lys Gly Glu Arg Leu Pro Gln Pro Pro Ile Cys Thr Ile Asp Val Tyr  
 930 935 940  
 Met Ile Met Val Lys Cys Trp Met Ile Asp Ala Asp Ser Arg Pro Lys  
 945 950 955 960  
 35 Phe Arg Glu Leu Ile Ile Glu Phe Ser Lys Met Ala Arg Asp Pro Gln  
 965 970 975  
 Arg Tyr Leu Val Ile Gln Gly Asp Glu Arg Met His Leu Pro Ser Pro  
 980 985 990  
 Thr Asp Ser Asn Phe Tyr Arg Ala Leu Met Asp Glu Glu Asp Met Asp  
 995 1000 1005  
 45 Asp Val Val Asp Ala Asp Glu Tyr Leu Ile Pro Gln Gln Gly Phe Phe  
 1010 1015 1020  
 Ser Ser Pro Ser Thr Ser Arg Thr Pro Leu Leu Ser Ser Leu Ser Ala  
 1025 1030 1035 1040  
 50 Thr Ser Asn Asn Ser Thr Val Ala Cys Ile Asp Arg Asn Gly Leu Gln  
 1045 1050 1055

55



S r Cys Pro Ile Lys Glu Asp Ser Phe Leu Gln Arg Tyr Ser Ser Asp  
 1060 1065 1070  
 6 Pro Thr Gly Ala Leu Thr Glu Asp Ser Ile Asp Asp Thr Phe Leu Pro  
 1075 1080 1085  
 Val Pro Glu Tyr Ile Asn Gln Ser Val Pro Lys Arg Pro Ala Gly Ser  
 1090 1095 1100  
 10 Val Gln Asn Pro Val Tyr His Asn Gln Pro Leu Asn Pro Ala Pro Ser  
 1105 1110 1115 1120  
 Arg Asp Pro His Tyr Gln Asp Pro His Ser Thr Ala Val Gly Asn Pro  
 1125 1130 1135  
 15 Glu Tyr Leu Asn Thr Val Gln Pro Thr Cys Val Asn Ser Thr Phe Asp  
 1140 1145 1150  
 Ser Pro Ala His Trp Ala Gln Lys Gly Ser His Gln Ile Ser Leu Asp  
 1155 1160 1165  
 20 Asn Pro Asp Tyr Gln Gln Asp Phe Phe Pro Lys Glu Ala Lys Pro Asn  
 1170 1175 1180  
 Gly Ile Phe Lys Gly Ser Thr Ala Glu Asn Ala Glu Tyr Leu Arg Val  
 1185 1190 1195 1200  
 25 Ala Pro Gln Ser Ser Glu Phe Ile Gly Ala  
 1205 1210

## (2) INFORMATION FOR SEQ ID NO:8:

30 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 1255 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: unknown  
 (D) TOPOLOGY: unknown

35 (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

40 Met Glu Leu Ala Ala Leu Cys Arg Trp Gly Leu Leu Leu Ala Leu Leu  
 1 5 10 15  
 Pro Pro Gly Ala Ala Ser Thr Gln Val Cys Thr Gly Thr Asp Met Lys  
 20 25 30  
 45 Leu Arg Leu Pro Ala Ser Pro Glu Thr His Leu Asp Met Leu Arg His  
 35 40 45  
 Leu Tyr Gln Gly Cys Gln Val Val Gln Gly Asn Leu Glu Leu Thr Tyr  
 50 55 60

50

55

Leu Pro Thr Asn Ala Ser Leu Ser Phe Leu Gln Asp Ile Gln Glu Val  
 65 70 75 80  
 5 Gln Gly Tyr Val Leu Ile Ala His Asn Gln Val Arg Gln Val Pro Leu  
 85 90 95  
 Gln Arg Leu Arg Ile Val Arg Gly Thr Gln Leu Phe Glu Asp Asn Tyr  
 100 105 110  
 10 Ala Leu Ala Val Leu Asp Asn Gly Asp Pro Leu Asn Asn Thr Thr Pro  
 115 120 125  
 Val Thr Gly Ala Ser Pro Gly Gly Leu Arg Glu Leu Gln Leu Arg Ser  
 130 135 140  
 15 Leu Thr Glu Ile Leu Lys Gly Gly Val Leu Ile Gln Arg Asn Pro Gln  
 145 150 155 160  
 Leu Cys Tyr Gln Asp Thr Ile Leu Trp Lys Asp Ile Phe His Lys Asn  
 165 170 175  
 20 Asn Gln Leu Ala Leu Thr Leu Ile Asp Thr Asn Arg Ser Arg Ala Cys  
 180 185 190  
 His Pro Cys Ser Pro Met Cys Lys Gly Ser Arg Cys Trp Gly Glu Ser  
 195 200 205  
 25 Ser Glu Asp Cys Gln Ser Leu Thr Arg Thr Val Cys Ala Gly Gly Cys  
 210 215 220  
 Ala Arg Cys Lys Gly Pro Leu Pro Thr Asp Cys Cys His Glu Gln Cys  
 225 230 235 240  
 30 Ala Ala Gly Cys Thr Gly Pro Lys His Ser Asp Cys Leu Ala Cys Leu  
 245 250 255  
 His Phe Asn His Ser Gly Ile Cys Glu Leu His Cys Pro Ala Leu Val  
 260 265 270  
 35 Thr Tyr Asn Thr Asp Thr Phe Glu Ser Met Pro Asn Pro Glu Gly Arg  
 275 280 285  
 Tyr Thr Phe Gly Ala Ser Cys Val Thr Ala Cys Pro Tyr Asn Tyr Leu  
 290 295 300  
 40 Ser Thr Asp Val Gly Ser Cys Thr Leu Val Cys Pro Leu His Asn Gln  
 305 310 315 320  
 45 Glu Val Thr Ala Glu Asp Gly Thr Gln Arg Cys Glu Lys Cys Ser Lys  
 325 330 335  
 Pro Cys Ala Arg Val Cys Tyr Gly Leu Gly Met Glu His Leu Arg Glu  
 340 345 350  
 50 Val Arg Ala Val Thr Ser Ala Asn Ile Gln Glu Phe Ala Gly Cys Lys  
 355 360 365

55

Lys Ile Phe Gly Ser Leu Ala Phe Leu Pr Glu Ser Phe Asp Gly Asp  
 370 375 380  
 5 Pro Ala Ser Asn Thr Ala Pro Leu Gln Pro Glu Gln Leu Gln Val Phe  
 385 390 395 400  
 Glu Thr Leu Glu Glu Ile Thr Gly Tyr Leu Tyr Ile Ser Ala Trp Pr  
 405 410 415  
 10 Asp Ser Leu Pro Asp Leu Ser Val Phe Gln Asn Leu Gln Val Ile Arg  
 420 425 430  
 Gly Arg Ile Leu His Asn Gly Ala Tyr Ser Leu Thr Leu Gln Gly Leu  
 435 440 445  
 15 Gly Ile Ser Trp Leu Gly Leu Arg Ser Leu Arg Glu Leu Gly Ser Gly  
 450 455 460  
 Leu Ala Leu Ile His His Asn Thr His Leu Cys Phe Val His Thr Val  
 465 470 475 480  
 20 Pro Trp Asp Gln Leu Phe Arg Asn Pro His Gln Ala Leu Leu His Thr  
 485 490 495  
 Ala Asn Arg Pro Glu Asp Glu Cys Val Gly Glu Gly Leu Ala Cys His  
 500 505 510  
 25 Gln Leu Cys Ala Arg Arg Ala Leu Leu Gly Ser Gly Pro Thr Gln Cys  
 515 520 525  
 Val Asn Cys Ser Gln Phe Leu Arg Gly Gln Glu Cys Val Glu Glu Cys  
 530 535 540  
 30 Arg Val Leu Gln Gly Leu Pro Arg Glu Tyr Val Asn Ala Arg His Cys  
 545 550 555 560  
 Leu Pro Cys His Pro Glu Cys Gln Pro Gln Asn Gly Ser Val Thr Cys  
 565 570 575  
 35 Phe Gly Pro Glu Ala Asp Gln Cys Val Ala Cys Ala His Tyr Lys Asp  
 580 585 590  
 Pro Pro Phe Cys Val Ala Arg Cys Pro Ser Gly Val Lys Pro Asp Leu  
 595 600 605  
 40 Ser Tyr Met Pro Ile Trp Lys Phe Pro Asp Glu Glu Gly Ala Cys Gln  
 610 615 620  
 Pro Cys Pro Ile Asn Cys Thr His Ser Cys Val Asp Leu Asp Asp Lys  
 625 630 635 640  
 45 Gly Cys Pro Ala Glu Gln Arg Ala Ser Pro Leu Thr Ser Ile Val Ser  
 645 650 655  
 Ala Val Val Gly Ile Leu Leu Val Val Val Leu Gly Val Val Phe Gly  
 660 665 670  
 50

55

Ile Leu Ile Lys Arg Arg Gln Gln Lys Ile Arg Lys Tyr Thr Met Arg  
 675 680 685  
 5 Arg Leu Leu Gln Glu Thr Glu Leu Val Glu Pro Leu Thr Pro Ser Gly  
 690 695 700  
 Ala Met Pro Asn Gln Ala Gln Met Arg Ile Leu Lys Glu Thr Glu Leu  
 705 710 715 720  
 10 Arg Lys Val Lys Val Leu Gly Ser Gly Ala Phe Gly Thr Val Tyr Lys  
 725 730 735  
 Gly Ile Trp Ile Pro Asp Gly Glu Asn Val Lys Ile Pro Val Ala Ile  
 740 745 750  
 15 Lys Val Leu Arg Glu Asn Thr Ser Pro Lys Ala Asn Lys Glu Ile Leu  
 755 760 765  
 Asp Glu Ala Tyr Val Met Ala Gly Val Gly Ser Pro Tyr Val Ser Arg  
 770 775 780  
 20 Leu Leu Gly Ile Cys Leu Thr Ser Thr Val Gln Leu Val Thr Gln Leu  
 785 790 795 800  
 Met Pro Tyr Gly Cys Leu Leu Asp His Val Arg Glu Asn Arg Gly Arg  
 805 810 815  
 25 Leu Gly Ser Gln Asp Leu Leu Asn Trp Cys Met Gln Ile Ala Lys Gly  
 820 825 830  
 Met Ser Tyr Leu Glu Asp Val Arg Leu Val His Arg Asp Leu Ala Ala  
 835 840 845  
 30 Arg Asn Val Leu Val Lys Ser Pro Asn His Val Lys Ile Thr Asp Phe  
 850 855 860  
 Gly Leu Ala Arg Leu Leu Asp Ile Asp Glu Thr Glu Tyr His Ala Asp  
 865 870 875 880  
 35 Gly Gly Lys Val Pro Ile Lys Trp Met Ala Leu Glu Ser Ile Leu Arg  
 885 890 895  
 Arg Arg Phe Thr His Gln Ser Asp Val Trp Ser Tyr Gly Val Thr Val  
 900 905 910  
 40 Trp Glu Leu Met Thr Phe Gly Ala Lys Pro Tyr Asp Gly Ile Pro Ala  
 915 920 925  
 Arg Glu Ile Pro Asp Leu Leu Glu Lys Gly Glu Arg Leu Pro Gln Pro  
 930 935 940  
 45 Pro Ile Cys Thr Ile Asp Val Tyr Met Ile Met Val Lys Cys Trp Met  
 945 950 955 960  
 50 Ile Asp Ser Glu Cys Arg Pro Arg Phe Arg Glu Leu Val Ser Glu Phe  
 965 970 975

55

Ser Arg Met Ala Arg Asp Pro Gln Arg Phe Val Val Ile Gln Asn Glu  
 980 985 990  
 6 Asp Leu Gly Pro Ala Ser Pro Leu Asp Ser Thr Phe Tyr Arg Ser Leu  
 995 1000 1005  
 Leu Glu Asp Asp Asp Met Gly Asp Leu Val Asp Ala Glu Glu Tyr Leu  
 1010 1015 1020  
 10 Val Pro Gln Gln Gly Phe Phe Cys Pro Asp Pro Ala Pro Gly Ala Gly  
 1025 1030 1035 1040  
 Gly Met Val His His Arg His Arg Ser Ser Ser Thr Arg Ser Gly Gly  
 1045 1050 1055  
 15 Gly Asp Leu Thr Leu Gly Leu Glu Pro Ser Glu Glu Glu Ala Pro Arg  
 1060 1065 1070  
 Ser Pro Leu Ala Pro Ser Glu Gly Ala Gly Ser Asp Val Phe Asp Gly  
 1075 1080 1085  
 20 Asp Leu Gly Met Gly Ala Ala Lys Gly Leu Gln Ser Leu Pro Thr His  
 1090 1095 1100  
 Asp Pro Ser Pro Leu Gln Arg Tyr Ser Glu Asp Pro Thr Val Pro Leu  
 1105 1110 1115 1120  
 25 Pro Ser Glu Thr Asp Gly Tyr Val Ala Pro Leu Thr Cys Ser Pro Gln  
 1125 1130 1135  
 Pro Glu Tyr Val Asn Gln Pro Asp Val Arg Pro Gln Pro Pro Ser Pro  
 1140 1145 1150  
 30 Arg Glu Gly Pro Leu Pro Ala Ala Arg Pro Ala Gly Ala Thr Leu Glu  
 1155 1160 1165  
 Arg Ala Lys Thr Leu Ser Pro Gly Lys Asn Gly Val Val Lys Asp Val  
 1170 1175 1180  
 35 Phe Ala Phe Gly Gly Ala Val Glu Asn Pro Glu Tyr Leu Thr Pro Gln  
 1185 1190 1195 1200  
 Gly Gly Ala Ala Pro Gln Pro His Pro Pro Pro Ala Phe Ser Pro Ala  
 1205 1210 1215  
 40 Phe Asp Asn Leu Tyr Tyr Trp Asp Gln Asp Pro Pro Glu Arg Gly Ala  
 1220 1225 1230  
 Pro Pro Ser Thr Phe Lys Gly Thr Pro Thr Val Ala Glu Asn Pro Glu  
 1235 1240 1245  
 45 Tyr Gly Leu Asp Val Pro Val  
 1250 1255

(2) INFORMATION FOR SEQ ID NO:9:

50

(i) SEQUENCE CHARACTERISTICS:

55

(A) LENGTH: 1342 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: unknown  
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

```

Met Arg Ala Asn Asp Ala Leu Gln Val Leu Gly Leu Leu Phe Ser Leu
1      5      10      15
Ala Arg Gly Ser Glu Val Gly Asn Ser Gln Ala Val Cys Pro Gly Thr
15     20     25     30
Leu Asn Gly Leu Ser Val Thr Gly Asp Ala Glu Asn Gln Tyr Gln Thr
35     40     45
Leu Tyr Lys Leu Tyr Glu Arg Cys Glu Val Val Met Gly Asn Leu Glu
20     50     55     60
Ile Val Leu Thr Gly His Asn Ala Asp Leu Ser Phe Leu Gln Trp Ile
65     70     75     80
Arg Glu Val Thr Gly Tyr Val Leu Val Ala Met Asn Glu Phe Ser Thr
25     85     90     95
Leu Pro Leu Pro Asn Leu Arg Val Val Arg Gly Thr Gln Val Tyr Asp
100    105    110
Gly Lys Phe Ala Ile Phe Val Met Leu Asn Tyr Asn Thr Asn Ser Ser
30     115    120    125
His Ala Leu Arg Gln Leu Arg Leu Thr Gln Leu Thr Glu Ile Leu Ser
35     130    135    140
Gly Gly Val Tyr Ile Glu Lys Asn Asp Lys Leu Cys His Met Asp Thr
145    150    155    160
Ile Asp Trp Arg Asp Ile Val Arg Asp Arg Asp Ala Glu Ile Val Val
40     165    170    175
Lys Asp Asn Gly Arg Ser Cys Pro Pro Cys His Glu Val Cys Lys Gly
180    185    190
Arg Cys Trp Gly Pro Gly Ser Glu Asp Cys Gln Thr Leu Thr Lys Thr
45     195    200    205
Ile Cys Ala Pro Gln Cys Asn Gly His Cys Phe Gly Pro Asn Pro Asn
210    215    220
Gln Cys Cys His Asp Glu Cys Ala Gly Gly Cys Ser Gly Pro Gln Asp
50     225    230    235    240

```

Thr Asp Cys Phe Ala Cys Arg His Phe Asn Asp Ser Gly Ala Cys Val  
 245 250 255  
 5 Pro Arg Cys Pro Gln Pro Leu Val Tyr Asn Lys Leu Thr Phe Gln Leu  
 260 265 270  
 Glu Pro Asn Pro His Thr Lys Tyr Gln Tyr Gly Gly Val Cys Val Ala  
 275 280 285  
 10 Ser Cys Pro His Asn Phe Val Val Asp Gln Thr Ser Cys Val Arg Ala  
 290 295 300  
 Cys Pro Pro Asp Lys Met Glu Val Asp Lys Asn Gly Leu Lys Met Cys  
 305 310 315 320  
 15 Glu Pro Cys Gly Gly Leu Cys Pro Lys Ala Cys Glu Gly Thr Gly Ser  
 325 330 335  
 Gly Ser Arg Phe Gln Thr Val Asp Ser Ser Asn Ile Asp Gly Phe Val  
 340 345 350  
 20 Asn Cys Thr Lys Ile Leu Gly Asn Leu Asp Phe Leu Ile Thr Gly Leu  
 355 360 365  
 Asn Gly Asp Pro Trp His Lys Ile Pro Ala Leu Asp Pro Glu Lys Leu  
 370 375 380  
 25 Asn Val Phe Arg Thr Val Arg Glu Ile Thr Gly Tyr Leu Asn Ile Gln  
 385 390 395 400  
 Ser Trp Pro Pro His Met His Asn Phe Ser Val Phe Ser Asn Leu Thr  
 405 410 415  
 30 Thr Ile Gly Gly Arg Ser Leu Tyr Asn Arg Gly Phe Ser Leu Leu Ile  
 420 425 430  
 Met Lys Asn Leu Asn Val Thr Ser Leu Gly Phe Arg Ser Leu Lys Glu  
 435 440 445  
 35 Ile Ser Ala Gly Arg Ile Tyr Ile Ser Ala Asn Arg Gln Leu Cys Tyr  
 450 455 460  
 His His Ser Leu Asn Trp Thr Lys Val Leu Arg Gly Pro Thr Glu Glu  
 465 470 475 480  
 Arg Leu Asp Ile Lys His Asn Arg Pro Arg Arg Asp Cys Val Ala Glu  
 485 490 495  
 40 Gly Lys Val Cys Asp Pro Leu Cys Ser Ser Gly Gly Cys Trp Gly Pro  
 500 505 510  
 Gly Pro Gly Gln Cys Leu Ser Cys Arg Asn Tyr Ser Arg Gly Gly Val  
 515 520 525  
 45 Cys Val Thr His Cys Asn Phe Leu Asn Gly Glu Pro Arg Glu Phe Ala  
 530 535 540  
 50

55

His Glu Ala Glu Cys Phe Ser Cys His Pro Glu Cys Gln Pro Met Gly  
 545 550 555 560  
 5 Gly Thr Ala Thr Cys Asn Gly Ser Gly Ser Asp Thr Cys Ala Gln Cys  
 565 570 575  
 Ala His Phe Arg Asp Gly Pro His Cys Val Ser Ser Cys Pro His Gly  
 580 585 590  
 10 Val Leu Gly Ala Lys Gly Pro Ile Tyr Lys Tyr Pro Asp Val Gln Asn  
 595 600 605  
 Glu Cys Arg Pro Cys His Glu Asn Cys Thr Gln Gly Cys Lys Gly Pro  
 610 615 620  
 15 Glu Leu Gln Asp Cys Leu Gly Gln Thr Leu Val Leu Ile Gly Lys Thr  
 625 630 635 640  
 His Leu Thr Met Ala Leu Thr Val Ile Ala Gly Leu Val Val Ile Phe  
 645 650 655  
 20 Met Met Leu Gly Gly Thr Phe Leu Tyr Trp Arg Gly Arg Arg Ile Gln  
 660 665 670  
 Asn Lys Arg Ala Met Arg Arg Tyr Leu Glu Arg Gly Glu Ser Ile Glu  
 675 680 685  
 25 Pro Leu Asp Pro Ser Glu Lys Ala Asn Lys Val Leu Ala Arg Ile Phe  
 690 695 700  
 Lys Glu Thr Glu Leu Arg Lys Leu Lys Val Leu Gly Ser Gly Val Phe  
 705 710 715 720  
 30 Gly Thr Val His Lys Gly Val Trp Ile Pro Glu Gly Glu Ser Ile Lys  
 725 730 735  
 Ile Pro Val Cys Ile Lys Val Ile Glu Asp Lys Ser Gly Arg Gln Ser  
 740 745 750  
 35 Phe Gln Ala Val Thr Asp His Met Leu Ala Ile Gly Ser Leu Asp His  
 755 760 765  
 Ala His Ile Val Arg Leu Leu Gly Leu Cys Pro Gly Ser Ser Leu Gln  
 770 775 780  
 Leu Val Thr Gln Tyr Leu Pro Leu Gly Ser Leu Leu Asp His Val Arg  
 785 790 795 800  
 45 Gln His Arg Gly Ala Leu Gly Pro Gln Leu Leu Leu Asn Trp Gly Val  
 805 810 815  
 Gln Ile Ala Lys Gly Met Tyr Tyr Leu Glu Glu His Gly Met Val His  
 820 825 830  
 50 Arg Asn Leu Ala Ala Arg Asn Val Leu Leu Lys Ser Pro Ser Gln Val  
 835 840 845

55



Gln Val Ala Asp Phe Gly Val Ala Asp Leu Leu Pro Pro Asp Asp Lys  
 850 855 860  
 6 Gln Leu Leu Tyr Ser Glu Ala Lys Thr Pro Ile Lys Trp Met Ala Leu  
 865 870 875 880  
 Glu Ser Ile His Phe Gly Lys Tyr Thr His Gln Ser Asp Val Trp Ser  
 885 890 895  
 10 Tyr Gly Val Thr Val Trp Glu Leu Met Thr Phe Gly Ala Glu Pro Tyr  
 900 905 910  
 Ala Gly Leu Arg Leu Ala Glu Val Pro Asp Leu Leu Glu Lys Gly Glu  
 915 920 925  
 15 Arg Leu Ala Gln Pro Gln Ile Cys Thr Ile Asp Val Tyr Met Val Met  
 930 935 940  
 Val Lys Cys Trp Met Ile Asp Glu Asn Ile Arg Pro Thr Phe Lys Glu  
 945 950 955 960  
 20 Leu Ala Asn Glu Phe Thr Arg Met Ala Arg Asp Pro Pro Arg Tyr Leu  
 965 970 975  
 Val Ile Lys Arg Glu Ser Gly Pro Gly Ile Ala Pro Gly Pro Glu Pro  
 980 985 990  
 25 His Gly Leu Thr Asn Lys Lys Leu Glu Glu Val Glu Leu Glu Pro Glu  
 995 1000 1005  
 Leu Asp Leu Asp Leu Asp Leu Glu Ala Glu Glu Asp Asn Leu Ala Thr  
 1010 1015 1020  
 30 Thr Thr Leu Gly Ser Ala Leu Ser Leu Pro Val Gly Thr Leu Asn Arg  
 1025 1030 1035 1040  
 Pro Arg Gly Ser Gln Ser Leu Leu Ser Pro Ser Ser Gly Tyr Met Pro  
 1045 1050 1055  
 35 Met Asn Gln Gly Asn Leu Gly Gly Ser Cys Gln Glu Ser Ala Val Ser  
 1060 1065 1070  
 Gly Ser Ser Glu Arg Cys Pro Arg Pro Val Ser Leu His Pro Met Pro  
 1075 1080 1085  
 40 Arg Gly Cys Leu Ala Ser Glu Ser Ser Glu Gly His Val Thr Gly Ser  
 1090 1095 1100  
 Glu Ala Glu Leu Gln Glu Lys Val Ser Met Cys Arg Ser Arg Ser Arg  
 1105 1110 1115 1120  
 45 Ser Arg Ser Pro Arg Pro Arg Gly Asp Ser Ala Tyr His Ser Gln Arg  
 1125 1130 1135  
 50 His Ser Leu Leu Thr Pro Val Thr Pro Leu Ser Pro Pro Gly Leu Glu  
 1140 1145 1150

55

Glu Glu Asp Val Asn Gly Tyr Val Met Pro A p Thr His Leu Lys Gly  
 1155 1160 1165  
 Thr Pro Ser Ser Arg Glu Gly Thr Leu Ser Ser Val Gly Leu Ser Ser  
 1170 1175 1180  
 Val Leu Gly Thr Glu Glu Glu Asp Glu Asp Glu Glu Tyr Glu Tyr Met  
 1185 1190 1195 1200  
 Asn Arg Arg Arg Arg His Ser Pro Pro His Pro Pro Arg Pro Ser Ser  
 1205 1210 1215  
 Leu Glu Glu Leu Gly Tyr Glu Tyr Met Asp Val Gly Ser Asp Leu Ser  
 1220 1225 1230  
 Ala Ser Leu Gly Ser Thr Gln Ser Cys Pro Leu His Pro Val Pro Ile  
 1235 1240 1245  
 Met Pro Thr Ala Gly Thr Thr Pro Asp Glu Asp Tyr Glu Tyr Met Asn  
 1250 1255 1260  
 Arg Gln Arg Asp Gly Gly Gly Pro Gly Gly Asp Tyr Ala Ala Met Gly  
 1265 1270 1275 1280  
 Ala Cys Pro Ala Ser Glu Gln Gly Tyr Glu Glu Met Arg Ala Phe Gln  
 1285 1290 1295  
 Gly Pro Gly His Gln Ala Pro His Val His Tyr Ala Arg Leu Lys Thr  
 1300 1305 1310  
 Leu Arg Ser Leu Glu Ala Thr Asp Ser Ala Phe Asp Asn Pro Asp Tyr  
 1315 1320 1325  
 Trp His Ser Arg Leu Phe Pro Lys Ala Asn Ala Gln Arg Thr  
 1330 1335 1340

## (2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 911 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: unknown  
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Met Lys Pro Ala Thr Gly Leu Trp Val Trp Val Ser Leu Leu Val Ala  
 1 5 10 15  
 Ala Gly Thr Val Gln Pro Ser Asp Ser Gln Ser Val Cys Ala Gly Thr  
 20 25 30

Glu Asn Lys Leu Ser Ser Leu Ser Asp Leu Glu Gln Gln Tyr Arg Ala  
 35 40 45  
 6 Leu Arg Lys Tyr Tyr Glu Asn Cys Glu Val Val Met Gly Asn Leu Glu  
 50 55 60  
 Ile Thr Ser Ile Glu His Asn Arg Asp Leu Ser Phe Leu Arg Ser Val  
 65 70 75 80  
 10 Arg Glu Val Thr Gly Tyr Val Leu Val Ala Leu Asn Gln Phe Arg Tyr  
 85 90 95  
 Leu Pro Leu Glu Asn Leu Arg Ile Ile Arg Gly Thr Lys Leu Tyr Glu  
 100 105 110  
 15 Asp Arg Tyr Ala Leu Ala Ile Phe Leu Asn Tyr Arg Lys Asp Gly Asn  
 115 120 125  
 Phe Gly Leu Gln Glu Leu Gly Leu Lys Asn Leu Thr Glu Ile Leu Asn  
 130 135 140  
 20 Gly Gly Val Tyr Val Asp Gln Asn Lys Phe Leu Cys Tyr Ala Asp Thr  
 145 150 155 160  
 Ile His Trp Gln Asp Ile Val Arg Asn Pro Trp Pro Ser Asn Leu Thr  
 165 170 175  
 25 Leu Val Ser Thr Asn Gly Ser Ser Gly Cys Gly Arg Cys His Lys Ser  
 180 185 190  
 Cys Thr Gly Arg Cys Trp Gly Pro Thr Glu Asn His Cys Gln Thr Leu  
 195 200 205  
 30 Thr Arg Thr Val Cys Ala Glu Gln Cys Asp Gly Arg Cys Tyr Gly Pro  
 210 215 220  
 Tyr Val Ser Asp Cys Cys His Arg Glu Cys Ala Gly Gly Cys Ser Gly  
 225 230 235 240  
 35 Pro Lys Asp Thr Asp Cys Phe Ala Cys Met Asn Phe Asn Asp Ser Gly  
 245 250 255  
 Ala Cys Val Thr Gln Cys Pro Gln Thr Phe Val Tyr Asn Pro Thr Thr  
 260 265 270  
 40 Phe Gln Leu Glu His Asn Phe Asn Ala Lys Tyr Thr Tyr Gly Ala Phe  
 275 280 285  
 Cys Val Lys Lys Cys Pro His Asn Phe Val Val Asp Ser Ser Ser Cys  
 290 295 300  
 45 Val Arg Ala Cys Pro Ser Ser Lys Met Glu Val Glu Glu Asn Gly Ile  
 305 310 315 320  
 50 Lys Met Cys Lys Pro Cys Thr Asp Ile Cys Pro Lys Ala Cys Asp Gly  
 325 330 335

55

Ile Gly Thr Gly Ser Leu Met Ser Ala Gln Thr Val Asp Ser Ser Asn  
 340 345 350  
 5 Ile Asp Lys Phe Ile Asn Cys Thr Lys Ile Asn Gly Asn Leu Ile Phe  
 355 360 365  
 Leu Val Thr Gly Ile His Gly Asp Pro Tyr Asn Ala Ile Glu Ala Ile  
 370 375 380  
 10 Asp Pro Glu Lys Leu Asn Val Phe Arg Thr Val Arg Glu Ile Thr Gly  
 385 390 395 400  
 Phe Leu Asn Ile Gln Ser Trp Pro Pro Asn Met Thr Asp Phe Ser Val  
 405 410 415  
 15 Phe Ser Asn Leu Val Thr Ile Gly Gly Arg Val Leu Tyr Ser Gly Leu  
 420 425 430  
 Ser Leu Leu Ile Leu Lys Gln Gln Gly Ile Thr Ser Leu Gln Phe Gln  
 435 440 445  
 20 Ser Leu Lys Glu Ile Ser Ala Gly Asn Ile Tyr Ile Thr Asp Asn Ser  
 450 455 460  
 Asn Leu Cys Tyr Tyr His Thr Ile Asn Trp Thr Thr Leu Phe Ser Thr  
 465 470 475 480  
 25 Ile Asn Gln Arg Ile Val Ile Arg Asp Asn Arg Lys Ala Glu Asn Cys  
 485 490 495  
 Thr Ala Glu Gly Met Val Cys Asn His Leu Cys Ser Ser Asp Gly Cys  
 500 505 510  
 30 Trp Gly Pro Gly Pro Asp Gln Cys Leu Ser Cys Arg Arg Phe Ser Arg  
 515 520 525  
 Gly Arg Ile Cys Ile Glu Ser Cys Asn Leu Tyr Asp Gly Glu Phe Arg  
 530 535 540  
 35 Glu Phe Glu Asn Gly Ser Ile Cys Val Glu Cys Asp Pro Gln Cys Glu  
 545 550 555 560  
 Lys Met Glu Asp Gly Leu Leu Thr Cys His Gly Pro Gly Pro Asp Asn  
 565 570 575  
 40 Cys Thr Lys Cys Ser His Phe Lys Asp Gly Pro Asn Cys Val Glu Lys  
 580 585 590  
 Cys Pro Asp Gly Leu Gln Gly Ala Asn Ser Phe Ile Phe Lys Tyr Ala  
 595 600 605  
 45 Asp Pro Asp Arg Glu Cys His Pro Cys His Pro Asn Cys Thr Gln Gly  
 610 615 620  
 Cys Asn Gly Pro Thr Ser His Asp Cys Ile Tyr Tyr Pro Trp Thr Gly  
 625 630 635 640  
 50  
 55

His Ser Thr Leu Pro Gln Asp Pro Val Lys Val Lys Ala Leu Glu Gly  
 645 650 655  
 6 Phe Pro Arg Leu Val Gly Pro Asp Phe Phe Gly Cys Ala Glu Pro Ala  
 660 665 670  
 Asn Thr Phe Leu Asp Pro Glu Glu Pro Lys Ser Cys Asp Lys Thr His  
 675 680 685  
 10 Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val  
 690 695 700  
 Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr  
 705 710 715 720  
 15 Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu  
 725 730 735  
 Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Val Ala Lys  
 740 745 750  
 20 Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser  
 755 760 765  
 Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys  
 770 775 780  
 25 Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile  
 785 790 795 800  
 Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro  
 805 810 815  
 30 Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu  
 820 825 830  
 Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn  
 835 840 845  
 35 Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser  
 850 855 860  
 Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg  
 865 870 875 880  
 40 Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu  
 885 890 895  
 His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
 900 905 910  
 45

## (2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 6 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: unknown

(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Gly Xaa Gly Xaa Xaa Gly  
1 5

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 6 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: unknown
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Asp Leu Ala Ala Arg Asn  
1 5

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 6 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: unknown
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Pro Ile Lys Trp Met Ala  
1 5

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:  
ACNGTNTGGG ARYTNAYHAC 20

8 (2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 23 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
10 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:  
CAYGTNAARA THACNGAYTT YGG 23

(2) INFORMATION FOR SEQ ID NO:16:

20 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 25 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: unknown

25 (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:  
30 GACGAATTCC NATHAARTGG ATGGC 25

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 24 base pairs  
(B) TYPE: nucleic acid  
35 (C) STRANDEDNESS: single  
(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:  
ACAYTTNARD ATDATCATRT ANAC 24

(2) INFORMATION FOR SEQ ID NO:18:

45 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 17 base pairs  
(B) TYPE: nucleic acid

50

55

- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

AANGTCATNA RYTCCCA

17

10

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 23 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: unknown

15

(ii) MOLECULE TYPE: DNA (genomic)

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

TCCAGNGCGA TCCAYTTDAT NGG

23

25

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 18 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: unknown

30

(ii) MOLECULE TYPE: DNA (genomic)

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

GGRTCDATCA TCCARCCT

18

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: unknown

40

(ii) MOLECULE TYPE: DNA (genomic)

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

50

55



CTGCTGTCAG CATCGATCAT

20

## (2) INFORMATION FOR SEQ ID NO:22:

- 5 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 7 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: unknown  
(D) TOPOLOGY: unknown

10 (ii) MOLECULE TYPE: peptide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

15 Thr Val Trp Glu Leu Met Thr  
1 5

## (2) INFORMATION FOR SEQ ID NO:23:

- 20 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 8 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: unknown  
(D) TOPOLOGY: unknown

25 (ii) MOLECULE TYPE: peptide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

30 His Val Lys Ile Thr Asp Phe Gly  
1 5

## (2) INFORMATION FOR SEQ ID NO:24:

- 35 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 7 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: unknown  
(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

40 Val Tyr Met Ile Ile Leu Lys  
1 5

## 45 (2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 6 amino acids

50

55

(B) TYPE: amino acid  
(C) STRANDEDNESS: unknown  
(D) TOPOLOGY: unknown

5

(ii) MOLECULE TYPE: peptid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

10

Trp Glu Leu Met Thr Phe  
1 5

(2) INFORMATION FOR SEQ ID NO:26:

15

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 8 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: unknown  
(D) TOPOLOGY: unknown

20

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

25

Pro Ile Lys Trp Met Ala Leu Glu  
1 5

(2) INFORMATION FOR SEQ ID NO:27:

30

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 6 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: unknown  
(D) TOPOLOGY: unknown

35

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

40

Cys Trp Met Ile Asp Pro  
1 5

(2) INFORMATION FOR SEQ ID NO:28:

45

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 35 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: unknown

50

(ii) MOLECULE TYPE: DNA (genomic)

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

GACTCGAGTC GACATCGATT TTTTTTTTTT TTTT

35

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

GAAGAAAGAC GACTCGTTCA TCGG

24

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

GACCATGACC ATGTAAACGT CAATA

25

# Claims

1. A recombinant polynucleotide comprising a sequence of at least about 200 nucleotides having greater than 80 % homology to a contiguous portion of the HER4 nucleotide sequence depicted in FIG. 1 or its complement.
2. The recombinant polynucleotide of claim 1 comprising a sequence of nucleotides encoding at least about 70 contiguous amino acids within the HER4 amino acid sequence depicted in FIG. 1.
3. The recombinant polynucleotide of claim 1 comprising a contiguous sequence of at least about 200 nucleotides within the HER4 nucleotide coding sequence depicted in FIG. 1 or its complement.
4. The recombinant polynucleotide of claim 1 comprising the HER4 nucleotide coding sequence depicted in FIG. 1 or its complement.
5. A recombinant polynucleotide which encodes a polypeptide having structural characteristics equivalent to that of HER4, which polynucleotide is obtained by single or multiple base addition, deletion and/or substitution in a nucleotide sequence of one of the claims 1 to 4, or which is obtained by selective hybridization with a nucleotide sequence of one of the claims 1 to 4.
6. A recombinant polynucleotide according to one of the claims 1 to 5 which is a DNA polynucleotide.
7. A recombinant polynucleotide according to one of the claims 1 to 5 which is a RNA polynucleotide.

8. An assay kit comprising a recombinant polynucleotide according to one of the claims 1 to 5 to which a detectable label has been added.
9. A polymerase chain reaction (PCR) kit comprising a pair of primers capable of priming cDNA synthesis in a PCR reaction, wherein each primer is a polynucleotide according to claim 6.
10. The PCR kit according to claim 9 further comprising a polynucleotide probe capable of hybridizing to a region of the HER4 gene between and not including the nucleotide sequences to which the primers hybridize.
11. A polypeptide comprising a sequence of at least about 80 amino acids having greater than 80 % identity to a contiguous portion of the HER4 amino acid sequence depicted in FIG. 1.
12. A HER4 polypeptide comprising
  - the amino acid sequence depicted in FIG. 1 from amino acid residues 1 through 1308; or
  - the amino acid sequence depicted in FIG. 1 from amino acid residues 26 through 1308; or
  - the amino acid sequence depicted in FIG. 1 from amino acid residues 1 through 1045; or
  - the amino acid sequence depicted in FIG. 1 from amino acid residues 26 through 1045; or
  - the amino acid sequence depicted in FIG. 2A; or
  - the amino acid sequence depicted in FIG. 1 from amino acid residues 772 through 1308; or
  - the amino acid sequence depicted in FIG. 2B.
13. A polypeptide having structural and/or functional features equivalent to HER4, obtainable by single or multiple amino acid addition, deletion and/or substitution in a sequence of one of the claims 11 or 12.
14. An antibody capable of inhibiting the interaction of a soluble polypeptide and human HER4.
15. An antibody according to claim 14 wherein the soluble polypeptide is a heregulin.
16. An antibody capable of
  - a) stimulating HER4 tyrosine autophosphorylation; or
  - b) inducing a HER4-mediated signal in a cell, which signal results in modulation of growth and/or differentiation of the cell; or
  - c) inhibiting HepG2 fraction 17-stimulated tyrosine phosphorylation of HER4 expressed in CHO/HER4 21-2 cells as deposited with the ATCC (accession number CRL 11205).
17. An antibody which immunospecifically binds to human HER4.
18. An antibody according to claim 17 which
  - a) resides on the cell surface after binding to HER4; or
  - b) is internalized into the cell after binding to HER4; or
  - c) immunospecifically binds to human HER4 expressed in CHO/HER4 21-2 cells as deposited with the ATCC (accession number CRL 11205); or
  - d) neutralizes HER4 biological activity; or
  - e) is conjugated to a drug or toxin; or
  - f) is radiolabeled.
19. Plasmid pBSHER4Y as deposited with the ATCC and having the accession number ATCC 69131.
20. A recombinant vector comprising a nucleotide sequence encoding a polypeptide according to one of the claims 11 to 13.
21. A host cell transfected with a recombinant vector according to claim 20.
22. A recombinant vector comprising a nucleotide sequence encoding a polypeptide according to one of the claims 11 to 13 wherein the coding sequence is operably linked to a control sequence which is capable of directing the expression of the coding sequence in a host cell transfected therewith.

23. A host cell transfected with a recombinant vector according to claim 22.
24. Cell line CHO/HER4 21-2 as deposited with the ATCC and having the accession number CRL 11205.
- 5 25. An assay for detecting the presence of a HER4 ligand in a sample comprising:  
(a) applying the sample to cells which have been engineered to overexpress HER4; and  
(b) detecting an ability of the ligand to affect an activity mediated by HER4.
- 10 26. The method according to claim 25, wherein the cells are CHO/HER4 21-2 cells as deposited with the ATCC and having the accession number CRL 11205.
27. The method according to claim 25, wherein the activity detected is HER4 tyrosine phosphorylation, or morphologic differentiation.
- 15 28. A ligand for HER4 comprising a polypeptide which binds to HER4, stimulates tyrosine phosphorylation of HER4, and affects a biological activity mediated by HER4.
29. A ligand according to claim 28 which is capable of inducing morphological differentiation when added to cultured MDA-MB-453 cells; and/or which is obtained from cultured HepG2 cell conditioned media.
- 20 30. An immunoassay for detecting HER4 comprising:  
a) providing an antibody according to claim 17 or 18;  
b) incubating a biological sample with the antibody under conditions which allow for the binding of the antibody to HER4; and  
25 c) determining the amount of antibody present as a HER4-antibody complex.
31. The use of at least one antibody according to one of the claims 17 or 18 for preparing a pharmaceutical composition for the in vivo delivery of a drug or toxin to cells expressing HER4.
- 30 32. The use of claim 31, which comprises conjugating at least one antibody according to claim 17 or 18, or an active fragment thereof, to the drug or toxin, for delivering the resulting conjugate to an individual by using a formulation, dose, and route of administration such that the conjugate binds to HER4.

35

40

45

50

55

1	MetLysProAlaThrGlyLeuTrpValTrpValSerLeuLeuValAlaAlaGlyThr	ValSerLeuLeuValAlaAlaGlyThr
1	ATTGTGACGACGGGATCTGAGACTTCACAAAA	GTGAGCCTTCTCTGCTGGCGCGGGGAC
20	ValGlnProSerAspSerGlnSerValCysAla	AspLeuGluGlnGlnTrpArgAlaLeu
91	GTCGAGCCAGCGATTCTCAGTCAGTGTGCGA	GACCTGGAAACAGCAGTACCGAGCCTTG
50	ArgLysTyrTyrGluAsnCysGluValValMet	AsnArgAspLeuSerPheLeuArgSer
181	CGCAAGTACTATGAAAACCTGTGAGGTGTGATG	AACCGGAGCTCTCTCTCTCTGCGGTCT
80	ValArgGluValThrGlyTyrValLeuValAla	AsnLeuArgIleIleArgGlyThrLys
271	GTTCGAGAAAGTCACAGGCTACGTGTTAGTGCT	AATTTAGCCATTATTCTGGGACAAAA
110	LeuTyrGluAspArgTyrAlaLeuAlaIlePhe	LeuGlnGluLeuGlyLeuLysAsnLeu
361	CTTTATGAGGATCGATATGCTTGGCAATATTT	CTTCAAGAACTTGGATTAAAGAACTTG
140	ThrGluIleLeuAsnGlyGlyValTyrValAsp	IleHisTyrPglAspIleValArgAsn
451	ACA GAAATCCTAAATGGTGCAGTCTATGTAGAC	ATTCAATGGCAAGATATTGTTCCGAAAC
170	ProTrpProSerAsnLeuThrLeuValSerThr	LysSerCysThrGlyArgCysTyrGly
541	CCA TGGCCTTCCAACTTGACTCTTGTGTCAACA	AAGTCTGTACTGGCCGTTGCTGGGGA
200	ProThrGluAsnHisCysGlnThrLeuThrArg	CysTyrGlyProTyrValSerAspCys
631	CCCACAGAAAATCATTGCACAGCTTTCACAAGG	TGCTACGGACCTTACGTCAGTGACTGC
230	CysHisArgGluCysAlaGlyGlyCysSerGly	AsnPheAsnAspSerGlyAlaCysVal
721	TGC CATCGAAATGTCTCGAGGCTGCTCAGGA	AATTTCAATGACACTGGAGCATGTGTT
260	ThrGlnCysProGlnThrPheValTyrAsnPro	AlaLysTyrThrTyrGlyAlaPheCys
811	ACTCAGTGTCCCCAAACCTTTGTCTACAATCCA	GCAAGTACACATATGGAGCACTTCTGT
290	ValLysLysCysProHisAsnPheValValAsp	SerLysMetGluValGluGluAsnGly
901	GTC AAGAATGTCCACATAACTTTGTGTAGAT	TCCAAGATGGAAGTAGAAGAAAAATGGG
320	IleLysMetCysLysProCysThrAspIleCys	SerLeuMetSerAlaGlnThrValAsp
991	ATT AAAATGTGTAAACCTTGCACCTGACATTTCG	TCAATGATGTAGCTCAGACTGTGGAT

### Figure 1

350 Ser SerAsnIleAspLysPheIleAsnCysThr LysIleAsnGlyAsnLeuIlePheLeuVal ThrGlyIleHisGlyAspProTyrAsn  
 1081 TCC AGTAACATTGACAAATTGATTAATGCTTCTAGTC AAGATCAATGGGAATTGATCTTCTAGTC ACTGGTATTCAATGGGACCTTACAAAT  
 380 Ala IleGluAlaIleAspProGluLysLeuAsn ValPheArgThrValArgGluIleThrGly PheLeuAsnIleGlnSerTyrProPro  
 1171 GCA ATTGAGCCATAGACCCAGAGAACTGAAC GTCTTTCCGACAGTCAAGAGATAACAGGT TTCCTGAACATACAGTCATGCGCCACCA  
 410 Asn MetThrAspPheSerValPheSerAsnLeu ValThrIleGlyGlyArgValLeuTyrSer GlyLeuSerLeuLeuIleLeuLysGln  
 1261 AAC ATGACTGACTTCAGTGTCTTTCTTAACCTG GTACCAATTGGTGAAGAGTACTCTATAGT GGCCTGTCTCTGCTTATCTCAAGCA  
 440 Gln GlyIleThrSerLeuGlnPheGlnSerLeu LysGluIleSerAlaGlyAsnIleTyrIle ThrAspAsnSerAsnLeuCysTyrTyr  
 1351 CAG GGCATCCTCTCTACAGTTCAGTCCCTG AAGGAATCAGCGCAGGAACATCTATATT ACTGACAAACAGCAACCTGTGTATTAT  
 470 His ThrIleAsnTyrThrThrLeuPheSerThr IleAsnGlnArgIleValIleArgAspAsn ArgLysAlaGluAsnCysThrAlaGlu  
 1441 CAT ACCATTAACTGGACACACTCTTCAGCACA ATCAACCAAGAGATAGTATATCGGGACAAAC AGAAAGCTGAAATTTGTACTGCTGAA  
 500 Gly MetValCysAsnHisLeuCysSerSerAsp GlyCysTyrGlyProGlyProAspGlnCys LeuSerCysArgArgPheSerArgGly  
 1531 GGA ATGGTGTGCAACCAATCTGTGTCCAGTGAT GGCTGTTGGGACCTGGGCCAATCCAAATGT CTGTCTGTCTGCGCCCTTCAGTAGAGGA  
 530 Arg IleCysIleGluSerCysAsnLeuTyrAsp GlyGluPheArgGluPheGluAsnGlySer IleCysValGluCysAspProGlnCys  
 1621 AGG ATCTGCATAGAGTCTTTGTAACCTCTATGAT GGTGAATTTGGGAGTTCAGAGATGGCTCC ATCTGTGTGGAGTGTGACCCCACTGT  
 560 Glu LysMetGluAspGlyLeuLeuThrCysHis GlyProGlyProAspAsnCysThrLysCys SerHisPheLysAspGlyProAsnCys  
 1711 GAG AAGATGGAAGATGGCTCTCTCACAATGCCAT GGACCGGCTCTGACAACCTGTACAAAGTGC TCTCATTTTAAAGATGGCCCAAACTGT  
 590 Val GluLysCysProAspGlyLeuGlnGlyAla AsnSerPheIlePheLysTyrAlaAspPro AspArgGluCysHisProCysHisPhe  
 1801 GTG GAAAAATGTCCAGATGGCTTACAGGGGGCA AACAGTTTTCATTTTCAAGTATGCTGATCCA GATCGGAGTGGCACCCTCATGCCATCCA  
 620 Asn CysThrGlnGlyCysAsnGlyProThrSer HisAspCysIleTyrTyrProTyrThrGly HisSerThrLeuProGlnHisAlaArg  
 1891 AAC TGCACCAAGGGTGAACGGTCCCACTAGT CATGACTGCATTTACTACCAATGGACGGGC CATTCACACTTTTACCACAAACATGCTAGA  
 650 Thr ProLeuIleAlaAlaGlyValIleGlyGly LeuPheIleLeuValIleValGlyLeuThr PheAlaValTyrValArgArgLysSer  
 1981 ACT CCCCTGATTGCAGCTGGAGTAATGGTGGG CTCTTCATTTCTGGTCAATTGTGGTCTGACA TTTGCTGTTTATGTTAGAGGAGAGC  
 680 Ile LysLysLysArgAlaLeuArgArgPheLeu GluThrGluLeuValGluProLeuThrPro SerGlyThrAlaProAsnGlnAlaGln  
 2071 ATC AAAAAAGAAAGACCTTGAGAGATTCTTG GAACAGAGTTGGTGGAAACCAATTAACTCCC AGTGGCACAGCACCCCAATCAAGCTCAA

Figure 1

(continued)

710 Leu ArgIleLeuLysGluThrGluLeuLysArg VallYsValLeuGlySerGlyAlaPheGly ThrValTyrLysGlyIleTyrValPro  
 2161 CTT CGTATTTTGAAGAACTGAGCTGAGAGG GTAAAGTCCTTGGTCAGGTGCTTTTGGG ACGTTTATAAAGGTATTTGGGTACCT  
 740 Glu GlyGluThrValLysIleProValAlaIle LysIleLeuAsnGluThrThrGlyProLys AlaAsnValGluPheMetAspGluAla  
 2251 GAA GGAGAAACTGTGAAGATTCTGTGCTATT AAGATTTCTTAATGACAACTGCTCCNAG GCAATGTGGAGTTTCATGGATGAAGCT  
 770 Leu IleMetAlaSerMetAspHisProHisLeu ValArgLeuLeuGlyValCysLeuSerPro ThrIleGlnLeuValThrGlnLeuMet  
 2341 CTG ATCATGGCAAGTATGGATCATCCACACCTA GTCCGGTTGCTGGGTGTGTCTGAGCCCA ACCATCCAGCTGGTACTCACTTATG  
 800 Pro HisGlyCysLeuLeuGluTyrValHisGlu HisLysAspAsnIleGlySerGlnLeuLeu LeuAsnTyrCysValGlnIleAlaLys  
 2431 CCC CATGGCTGCTGTGGAGTATGTCACGAG CACAAGGATAACATTGGATCACAACTGCTG CTTAACTGGTGTGTCTCCAGATAGCTAG  
 830 Gly MetMetTyrLeuGluGluArgArgLeuVal HisArgAspLeuAlaAlaArgAsnValLeu ValLysSerProAsnHisValLysIle  
 2521 GGA ATGATGTACCTGGAAGAAAGACGACTCGT CATCGGATTGGCAGCCCGTAATGCTTA GTGAAATCTCCAAACCATGTGAAATC  
 860 Thr AspPheGlyLeuAlaArgLeuGluGly AspGluLysGluTyrAsnAlaAspGlyGly LysMetProIleLysTyrMetAlaLeu  
 2611 ACA GATTTTGGCTAGCCAGACTCTTTGGAAGGA GATGAAGAGAGTACATGCTGATGGAGGA AAGATGCCAATTAATGGATGCTCTG  
 890 Glu CysIleHisTyrArgLysPheThrHisGln SerAspValTyrSerTyrGlyValThrIle TyrGluLeuMetThrPheGlyGlyLys  
 2701 GAG TGTATACATTACAGGAATTCACCCATCAG AGTGACGTTTGGAGCTATGGAGTACTATA TGGGAAGTATGACCTTTGGAGGAAAA  
 920 Pro TyrAspGlyIleProThrArgGluIlePro AspLeuLeuGluLysGlyGluArgLeuPro GlnProIleCysThrIleAspVal  
 2791 CCC TATGATGGAATTCACCGGAGAAATCCCT GATTTATAGAGAAAGGAGAACGTTTGCCCT CAGCTCCCATCTGCACATATGACGTT  
 950 Tyr MetValMetValLysCysTyrMetIleAsp AlaAspSerArgProLysPheLysGluLeu AlaAlaGluPheSerArgMetAlaArg  
 2881 TAC ATGGTCATGCTCAATGTTGGATGATTGAT CCTGACAGTAGACCTAAATTTAAGGAAGTCTGCTGAGTTTTTCAGGATGGCTCGA  
 980 Asp ProGlnArgTyrLeuValIleGlnGlyAsp AspArgMetLysLeuProSerProAsnAsp SerLysPhePheGlnAsnLeuLeuAsp  
 2971 GAC CCTCAAGATACCTAGTTATTCAGGGTGAT GATCGTATGAGCTTCCAGTCCAAATGAC AGCAAGTTCTTTTCAGAAATCTCTTGGAT  
 1010 Glu GluAspLeuGluAspMetMetAspAlaGlu GluTyrLeuValProGlnAlaPheAsnIle ProProIleTyrThrSerArgAla  
 3061 GAA GAGGATTTGCAAGATATGATGATGCTGAG GAGTACTTGGTCCCTCAGGCTTTCACATC CCACCTCCCATCTATATCTCCAGAGCA  
 1040 Arg IleAspSerAsnArgSerGluIleGlyHis SerProProProAlaTyrThrProMetSer GlyAsnGlnPheValTyrArgAspGly  
 3151 AGA ATTGACTCGAATAGGAGTGAATTTGGACAC AGCCCTCTCTCTGCTGCTACACCCCATGTCA GGAACCAAGTTTGTATATCCGAGATGGA  
 1070 Gly PheAlaAlaGluGlnGlyValSerValPro TyrArgAlaProThrSerThrIleProGlu AlaProValAlaGlnGlyAlaThrAla  
 3241 GGT TTTGCTGCTGAACACAGGAGTGTCTGTGCC TACAGAGCCCACTAGCACCAATTCAGAA GCTCTGTGGCACAGGGTCTACTGCT  
 1100 Glu IlePheAspAspSerCysAsnGlyThr LeuArgLysProValAlaProHisValGln GluAspSerSerThrGlnArgTyrSer  
 3331 GAG ATTTTGTGACTCTCTGCTGTATATGGCACC CTACGCAAGCCAGTGGCACCCCATGTCCAA GAGGACAGTAGACCCAGAGGTACAGT

Figure 1  
(continued)



Figure 1  
(continued)

### Figure 2A

350 Ser SerAsnIleAspLysPheIleAsnCysThr LysIleAsnGlyAsnLeuIlePheLeuVal ThrGlyIleHisGlyAspProTyrAsn  
 1081 TCC AGTAACATTGACAAATTCATAAAGTACC AAGATCAATGCGAATTGATCTTTCTAGTC ACTGGTATTTCATGGGACCCCTTACAAAT  
 380 Ala IleGluAlaIleAspProGluLysLeuAsn ValPheArgThrValArgGluIleThrGly PheLeuAsnIleGlnSerTyrProPro  
 1171 GCA ATTGAAGCCATAGACCCAGAGAACTGAAC GTCTTTCCGACAGTCAGAGAGAAACAGGT TTCTCTAACATACAGTCAATGGCCACCA  
 410 Asn MetThrAspPheSerValPheSerAsnLeu ValThrIleGlyGlyArgValLeuTyrSer GlyLeuSerLeuLeuIleLeuLysGln  
 1261 AAC ATGACTGACTTCAGTGTGTTTTTCTAACCTG GTGACCAATGCTGGAGAGTACTCTATAGT GGCCTGCTCTGCTTATCTCTCAAGCAA  
 440 Gln GlyIleThrSerLeuGlnPheGlnSerLeu LysGluIleSerAlaGlyAsnIleTyrIle ThrAspAsnSerAsnLeuCysTyrTyr  
 1351 CAG GGCATCACCTCTCTACAGTTCAGTCCCTG AAGGAATCAGCGCAGGAACATCTATATT ACTGACAAACAGCACTGTGTTATTAT  
 470 His ThrIleAsnTyrThrThrLeuPheSerThr IleAsnGlnArgIleValIleArgAspAsn ArgLysAlaGluAsnCysThrAlaGlu  
 1441 CAT ACCATTAACCTGGACAACACTCTGTGTTCCAGTGAT GGTGAATTTCCGGAGTTTGAGATGGCTCC ATCTGTGTGAGTGTCAACCCCACTGT  
 500 Gly MetValCysAsnHisLeuCysSerSerAsp GlyCysTyrGlyProGlyProAspGlnCys LeuSerCysArgArgPheSerArgGly  
 1531 GGA ATGGTGTGCAACCATCTGTGTTCCAGTGAT GGCTGTTGGGACCTGGGCCAGACCAATGT CTGTCTGTCTCGCCGCTTCAGTAGAGGA  
 530 Arg IleCysIleGluSerCysAsnLeuTyrAsp GlyGluPheArgGluPheGluAsnGlySer IleCysValGluCysAspProGlnCys  
 1621 AGG ATCTGCATAGAGTCTTGTAACCTCTATGAT GGTGAATTTCCGGAGTTTGAGATGGCTCC ATCTGTGTGAGTGTCAACCCCACTGT  
 560 Glu LysMetGluAspGlyLeuLeuThrCysHis GlyProGlyProAspAsnCysThrLysCys SerHisPheLysAspGlyProAsnCys  
 1711 GAG AAGATGGAGATGGCTCTCTACATGCCAT GGACCGGTCTCTGACAACTGTACAAAGTGC TCTCATTTTAAAGATGGCCCAAACTGT  
 590 Val GluLysCysProAspGlyLeuGlnGlyAla AsnSerPheIlePheLysTyrAlaAspPro AspArgGluCysHisProCysHisPro  
 1801 GTG GAAAAAATGTCCAGATGGCTTACAGGGGGCA AACAGTTTTCATTTTCAAGTATGCTGATCCA GATCGGAGTGGCCACCCATGCCATCCA  
 620 Asn CysThrGlnGlyCysAsnGlyProThrSer HisAspCysIleTyrTyrProTyrThrGly HisSerThrLeuProGlnHisAlaArg  
 1891 AAC TGCACCCCAAGGTGTACGGTCCCACTAGT CATGACTGCATTTACTACCCATGGACGGGC CATTCACATTTTACCAACATCTCTAGA  
 650 Thr ProLeuIleAlaAlaGlyValIleGlyGly LeuPheIleLeuValIleValGlyLeuThr PheAlaValTyrValArgArgLysSer  
 1981 ACT CCCCTGATTGCAGCTGGAGTAAATGGTGGG CTCCTTCATTTCTGGTCNTTGGGCTCTGACA TTTCCTGTTTATGTTAGAGGAAGAGC

Figure 2A  
(continued)

680 Ile LysLysLysArgAlaLeuArgArgPheLeu GluThrGluLeuValGluProLeuThrPro SerGlyThrAlaProAnglnAlaGln  
 2071 ATC AAAAAGAAAAGAGCCCTTGAGAGATTCTTG GAAACAGAGTTGGTGGACCATTAACCTCC AGTGGACAGCACCCCAATCAAGCTCAA  
 710 Leu ArgIleLeuLysGluThrGluLeuLysArg ValLysValLeuGlySerGlyAlaPheGly ThrValTyrLysGlyIleTrpValPr  
 2161 CTT CGTATTTTGAAGAACTGAGCTGAGAGGG GTAAAGTCCTTGGCTCAGGTGCTTTTGGC ACGTTTATTAAGGTATTTGGGTACT  
 740 Glu GlyGluThrValLysIleProValAlaIle LysIleLeuAsnGluThrThrGlyProLys AlaAsnValGluPheMetAspGluAla  
 2251 GAA GGAGAAACTGTGAAGATTCTCTGTGGCTATT AGATTCTTAATGAGACAACTGTCTCCAG GCNAATGTGGAGTTTCATGATGAAGCT  
 770 Leu IleMetAlaSerMetAspHisProHisLeu ValArgLeuLeuGlyValCysLeuSerPro ThrIleGlnLeuValThrGlnLeuMet  
 2341 CTG ATCATGGCAAGTATGGATCATCCACACTA GTCCGGTTGCTGGTGTGTGTCTGAGCCCA ACCATCCAGCTGGTTACTCAACTATG  
 800 Pro HisGlyCysLeuLeuGluTyrValHisGlu HisLysAspAsnIleGlySerGlnLeuLeu LeuAsnTrpCysValGlnIleAlaLys  
 2431 CCC CATGGCTGCTGTGGAGTATGTCCAGAG CACAAGGATAACATTGGATCACACTGCTG CTTAAGTGGTGTGTCCAGATAGCTAAG  
 830 Gly MetMetTyrLeuGluGluArgArgLeuVal HisArgAspLeuAlaAlaArgAsnValLeu ValLysSerProAspHisValLysIle  
 2521 GGA ATGATGTACCTGCAAGAAAGACGACTCGTT CATCGGATTTGGCAGCCCGTAAATGCTTA GTGAAATCTCCAAACCATGTGAAATC  
 860 Thr AspPheGlyLeuAlaArgLeuLeuGluGly AspGluLysGluTyrAsnAlaArgGlyGly LysMetProIleLysTrpMetAlaLeu  
 2611 ACA GATTTTGGGCTAGCCAGACTCTTTGGAAGCA GATGAAAAGAGTACAAATGCTGTGAGGGA AGATGCCCAATTAAATGGATGGCTCTG  
 890 Glu CysIleHisTyrArgLysPheThrHisGln SerAspValTrpSerTyrGlyValThrIle TrpGluLeuMetThrPheGlyGlyLys  
 2701 GAG TGTATACATTACAGGAATTCACCCATCAG AGTGACGTTTGGAGCTATGGAGTACTATA TGGGAAGTGTATGACCTTTGGAGGAAA  
 920 Pro TyrAspGlyIleProThrArgGluIlePro AspLeuLeuGluLysGlyGluArgLeuPro GlnProProIleCysThrIleAspVal  
 2791 CCC TATGATGGAATTCACGCGGAGAAATCCCT GATTATTAGAGAAAGAGAGAACGTTTGCCT CAGCCTCCCATCTGCATATTGACGTT  
 950 Tyr MetValMetValLysCysTrpMetIleAsp AlaAspSerArgProLysPheLysGluLeu AlaAlaGluPheSerArgMetAlaArg  
 2881 TAC ATGGTCATGGTCAATGTTGGATGATTGAT GCTGACAGTAGACCTAAATTTAAGGAACTG CCTGCTGAGTTTTCAGAGGATGGCTCGA  
 980 Asp ProGlnArgTyrLeuValIleGlnGlyAsp AspArgMetLysLeuProSerProAsnAsp SerLysPhePheGlnAsnLeuLeuAsp  
 2971 GAC CCTCAAGATACCTACTATTATTCAGGGTGAT GATCGTATGAGCTTCCAGTCGAAATGAC AGCAAGTTCTTTCAGATCTCTTGGAT  
 1010 Glu GluAspLeuGluAspMetMetAspAlaGlu GluTyrLeuValProGlnAlaPheAsnIle ProProProIleTyrThrSerArgAla  
 3061 GAA GAGGATTTGGAGATATGATGGATGCTGAG GACTACTTGGTCCCTCAGGCTTTCACACATC CCACCTCCCATCTATATACTTCCAGAGCA

Figure 2A  
(continued)

[illegible]

**Figure 2A**  
(continued)

HER4 N-terminal truncated with AP domain

```

1  CATTAGCTGCAATTGATCAAGTCACTGAGAGAAGGGCAACATTCATCCACAGATATAGTGGTATGGGAAAGCCCTCGATGTTGA
85  AATCTAGCTTCAAAAAGCCCTGCTCGAAATGTAGTTAATTGGATGAAGTGAAGAGATATAAACACAGAGAG GAAGCTCTGATC

1  MetAlaSerMetAspHisProHisLeuVal ArgLeuLeuGlyValCysLeuSerProThr IleGlnLeuValThrGlnLeuMetProHis
168  ATGGCAAGTATGATCATCCACACCTAGTC CGGTTCTGGGTGTGTCTGAGCCCAACC ATCCAGCTGGTTACTCAACTTATGCCCCAT

31  GlyCysLeuLeuGluTyrValHisGluHis LysAspAsnIleGlySerGlnLeuLeuLeu AsnTrpCysValGlnIleAlaLysGlyMet
258  GGCTGCCCTGTTGGAGTATGTCCACGACAC AAGGATAACATTTGGATCACTACTGCTGTTT AACTGGTGTGCCAGATAGCTAAGGGAATG

61  MetTyrLeuGluGluArgArgLeuValHis ArgAspLeuAlaAlaArgAsnValLeuVal LysSerProAsnHisValLysIleThrAsp
348  ATGTACCTGGAGAGAAAGACGACTCGTTTCAT CGGGAATTGGCAGCCCGTAATGTCTTAGTG AATCTCCAAACCATGTGAAATTCACAGAT

91  PheGlyLeuAlaArgLeuLeuGluGlyAsp GluLysGluTyrAsnAlaAspGlyGlyLys MetProIleLysTrpMetAlaLeuGluCys
438  TTTGGGCTAGCCAGACTCTTGAAGGAGAT GAAAAAGAGTACAAATGCTGATGGAGGAAG ATGCCAATAAATGAATGGCTCTCGAGTGT

121  IleHisTyrArgLysPheThrHisGlnSer AspValTrpSerTyrGlyValThrIleTrp GluLeuMetThrPheGlyGlyLysProTyr
528  ATACATTACAGGAAATTCACCCATCAGAGT GACGTTGGAGCTATGGAGTTACTAATAGG GAACGTGATGACCTTTTGGAGGAAACCCCTAT

151  AspGlyIleProThrArgGluIleProAsp LeuLeuGluLysGlyGluArgLeuProGln ProProIleCysThrIleAspValTyrMet
618  GATGGAATCCAAACCGGAGAAATCCCTGAT TTATTAGAGAAAGGAGAACGTTTGGCTCAG CCTCCCATCTGCACATATTACGCTTTTACATG

181  ValMetValLysCysTrpMetIleAspAla AspSerArgProLysPheLysGluLeuAla AlaGluPheSerArgMetAlaArgAspPro
708  GTCATGGTCAAAATGTTGGATGATTGATGCT GACAGTAGACCTAAATTTAAGGNACTGGCT GCTGAGTTTTCAGGATGGCTCGAGACCCCT

211  GlnArgTyrLeuValIleGlnGlyAspAsp ArgMetLysLeuProSerProAsnAspSer LysPhePheGlnAsnLeuLeuAspGluGlu
798  CAAAGATACCTAGTTATTTCAGGGTGATGAT CGTATGAAAGCTTCCCAGTCCAAATGACAGC AAGTTCTTTCAGAAATCTCTTGGATGAGAG

241  AspLeuGluAspMetMetAspAlaGluGlu TyrLeuValProGlnAlaPheAsnIlePro ProProIleTyrThrSerArgAlaArgIle
888  GATTTGGAAGATATGATGGATGCTGAGGAG TACTTGGTCCCTCAGGCTTCAACATCCCA CCTCCCATCTATACTTCCAGAGCAAGAATT

271  AspSerAsnArgSerGluIleGlyHisSer ProProAlaTyrThrProMetSerGly AsnGlnPheValTyrArgAspGlyGlyPhe
978  GACTCGAATAGGAGTGAATTTGGACACAGC CCTCCTCTCTGCTACACCCCATGTCCAGGA AACCAATTGTATACCGAGATGGAGTTT

301  AlaAlaGluGlnGlyValSerValProTyr ArgAlaProThrSerThrIleProGluAla ProValAlaGlnGlyAlaThrAlaGluIle
1068  GCTGCTGAACAAGGAGTGTCTGTGCTCCCTAC AGAGCCCCCACTAGCACAAATTCAGAGCT CCTGTGGCACAGGGTGTCTACTGCTGAGATT

```

Figure 2B

(continued)

HER4  
HER4 with alternate 3'-end without Autophosphorylation domain

MKPATGLMWWVSLVAAGTVQPSDSQSV CAGTENKLSLSDLEQQYRALRKYYENCEVVM	60
::	
MKPATGLMWWVSLVAAGTVQPSDSQSV CAGTENKLSLSDLEQQYRALRKYYENCEVVM	60
GNLEITSIEHNRDLSFLRSVREVTGYVLVALNQFRYLPLENLRIIRGTKLYEDRYALAI F	120
::	
GNLEITSIEHNRDLSFLRSVREVTGYVLVALNQFRYLPLENLRIIRGTKLYEDRYALAI F	120
LNRYKDGNGFLQELGLKNLTEILNGGVYVDQNKFLCYADTIHQDIVRNPMPSNLT LVST	180
::	
LNRYKDGNGFLQELGLKNLTEILNGGVYVDQNKFLCYADTIHQDIVRNPMPSNLT LVST	180
NGSSGGRCHKSCCTGRONGPTENHQQT LTRTVCAEQCDGRCYGPYVSDOCHRECAGGCSG	240
::	
NGSSGGRCHKSCCTGRONGPTENHQQT LTRTVCAEQCDGRCYGPYVSDOCHRECAGGCSG	240
PKOTDCFACHNFENDSGACVTQCPQT FVYNPTTFQLEHNFNAKYTYGAFVKKCPHNFVD	300
::	
PKOTDCFACHNFENDSGACVTQCPQT FVYNPTTFQLEHNFNAKYTYGAFVKKCPHNFVD	300
SSSCVRACPSSKMEVEENGIMCKPCTD ICPKACDGIGTGSLSAQTVDSNIDKFINCT	360
::	
SSSCVRACPSSKMEVEENGIMCKPCTD ICPKACDGIGTGSLSAQTVDSNIDKFINCT	360
KINGNLIFLVTGIHGDPYNAIEAIDPEK LNVFRTVREITGFLNIQSWPPNMTDFS VFSNL	420
::	
KINGNLIFLVTGIHGDPYNAIEAIDPEK LNVFRTVREITGFLNIQSWPPNMTDFS VFSNL	420
VTIGGRVLYSGLSL LILKQQGITS LQFQSLKEISAGNIYITDNSNLCYYHTINW TTFST	480
::	
VTIGGRVLYSGLSL LILKQQGITS LQFQSLKEISAGNIYITDNSNLCYYHTINW TTFST	480
INQRIVIRDNRKAENCTAEGHVCNHLCS SDGCGPGPDQCLSCRRFSRGRICIESCNLYD	540
::	
INQRIVIRDNRKAENCTAEGHVCNHLCS SDGCGPGPDQCLSCRRFSRGRICIESCNLYD	540
GEFREFEINGSICVECDPQCEKME DGLLTCHGPGPDNCTKCSHFKDGPN CVEKCPDGLQGA	600
::	
GEFREFEINGSICVECDPQCEKME DGLLTCHGPGPDNCTKCSHFKDGPN CVEKCPDGLQGA	600

Figure 3A



NSFI FKYADPDRECHPCHPNCTQGCGPTSHDCIYYPWTGHSTLPQHARTFLIAGVIGG	660
NSFI FKYADPDRECHPCHPNCTQGCGPTSHDCIYYPWTGHSTLPQHARTFLIAGVIGG	660
LFILVIVGLTFAVYVRRKSIKKRALRRFLETTELVEPLTPSGTAPNQAQLRIKTELKR	720
LFILVIVGLTFAVYVRRKSIKKRALRRFLETTELVEPLTPSGTAPNQAQLRIKTELKR	720
VKVLGSGAGFTVYKGIWVPEGETVKI PVAIKILNETTGPKANVEFMDEALIMASHDHPHL	780
VKVLGSGAGFTVYKGIWVPEGETVKI PVAIKILNETTGPKANVEFMDEALIMASHDHPHL	780
VRLGVCLSPITQLVTQLMPHGCLLEYVHEHKDNIGSQLLNMCVQIAKGMYLEERRLV	840
VRLGVCLSPITQLVTQLMPHGCLLEYVHEHKDNIGSQLLNMCVQIAKGMYLEERRLV	840
HRDLAARNVLVKSPNHVKITDFGLARLLEGDEKEYNADGGKMPKMALECIHYRKETHQ	900
HRDLAARNVLVKSPNHVKITDFGLARLLEGDEKEYNADGGKMPKMALECIHYRKETHQ	900
SDVWSYGVTIWELMTFGGKPYDGIPTREIPDLLEKGERLPQPPICTIDVYHVMVKCWMID	960
SDVWSYGVTIWELMTFGGKPYDGIPTREIPDLLEKGERLPQPPICTIDVYHVMVKCWMID	960
ADSRPKFKELAAEFMRMARDPQRYLVIQGDDRMKLPSPNDSKFFQNLLEEDLEDMMDAE	1020
ADSRPKFKELAAEFMRMARDPQRYLVIQGDDRMKLPSPNDSKFFQNLLEEDLEDMMDAE	1020
EYLVPOAFNIPPPITYTSRARIIDSNRSEIGHSPPPAYTPMSGNQFYVRDGGFAAEQGVSVF	1080
EYLVPOAFNIPPPITYTSRARIIDSNRVRNNYIHIS-YSF	1057
YRAPSTIPEAPVAQGATAEIFDDSCNGTLRKPVAPHVQEDSSTORYSADPTVFAPERS	1140
PRGELDEEGYMTPMRDKPKQEYLNPEENPFVSRKNGDLQALDNPEYHNASNGPPKAED	1200
EYVNEPLYLNTFANTLGKAEYLNKNNILSMPEKAKKAFDNPDYWNHSLPPRSTLQHPDYLO	1260
EYSTKYFYKQNGRIRPVAENPEYLSFSLKPGTVLPPPPYRHRNTVV	1308

Aligned 1058, Matches 1046, Mismatches 12, Score 132, Homology 98%

**Figure 3A**  
(continued)

HER4

HER4 N-terminal truncated with autophosphorylation domain

```

MKPATGLNVHVSLLVAAGTVQPSDSQSVCACTENKLSLSLEQQYRALRKYYENCEVVM      60
GNLEITSIEHNRDLSFLRSVREVTGYVLVALNQFRLPLENLRIRGTNLYEDRYALAI      120
LNRYKDGNGFGLQELGLNLTIELNGGVYVDQNKFLCYADTIHMQOIVRNPNPNSLTLVST    180
NGSSGCGRCHKSCITGRNGPTENHCQTLTRTVCAEQCGRCYGPYVSDCCHRECAGGCSG    240
PKDTDCFACNNFNDGACVTQCPQTFVYNPTTFQLEHNFNAKYTYGAFVCVKCPHNFVVD    300
SSSCVRACPSNMEVEENGIHCKPCTDICPKACDGIOTGSLMSAQTVDSNIDKFINCT      360
KINGNLI FLVTGIHGDPYNAIEAIDPEKLNVTFTVREITGFLNIQSWPPNMTDFSVFSNL    420
VTIGGRVLYSGLSLLILKQQGITSLQFQSLKEISAGNIYITDNSNLCTYHTINWTLFST    480
INQRIVIRDNRKAENCTAEGMVCHLCSGDCNGPGPDQCLSCRRFSRGRI CIESCNLYD     540
GEFREFEENGSI CVECDPQCEKMEGDLTCHGPGPDNCTKCSHFKDGPNCEKCPDGLQGA    600
NSFIKRYADPDRECHPCHPNTQGCNGPTSHDCIYYPNTGHSTLPQHARTFLIAGVIGG     660

VKVLGSGAGFTVYKGIWVPEGETVKIPVAIKILNETTGPKANVEFMDEALIMASHDHPHL    780
      ::::::::::::::
      EALIMASHDHPHL      13

VRLLGVCLSPTIQLVTQLMPHGCLLEYVHEHKDNIQSLLLLNMCVQIAKGMHYLEERRLV    840
::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::
VRLLGVCLSPTIQLVTQLMPHGCLLEYVHEHKDNIQSLLLLNMCVQIAKGMHYLEERRLV    73

HRDLAARNVLVKSPNHVKITDFGLARLLEGDEKEYNADGGKMPKMMALECIHYRKETHQ      900
::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::
HRDLAARNVLVKSPNHVKITDFGLARLLEGDEKEYNADGGKMPKMMALECIHYRKETHQ      133

SDVMSYGYTIWELMTFGGKPYDGIPTREIPDLLEKGERLPQPPICTIDVYHVMVKCWMID     960
::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::
SDVMSYGYTIWELMTFGGKPYDGIPTREIPDLLEKGERLPQPPICTIDVYHVMVKCWMID     193

ADSRPKFELAAEFMRHARDPQRYLVIQGDDRMKLPSPNDSKFFQNLLEDEEDLEDMMAE    1020
::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::
ADSRPKFELAAEFMRHARDPQRYLVIQGDDRMKLPSPNDSKFFQNLLEDEEDLEDMMAE     253

EYLVPOAFNIPPIIYTSRARIDSNRSEIGHSPPPAYTPMSGNQFVYRDGGFAAEQGVSV      1080
::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::
EYLVPOAFNIPPIIYTSRARIDSNRSEIGHSPPPAYTPMSGNQFVYRDGGFAAEQGVSV      313

YRAPSTSTIPEAPVAQGATAEIFDDSCNGTLRKFPVAPHVQEDSSTQRYSDPTVFAPERS    1140
::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::
YRAPSTSTIPEAPVAQGATAEIFDDSCNGTLRKFPVAPHVQEDSSTQRYSDPTVFAPERS    373

PRGELDEEGYTMPMRDKPKQEYLNPEENPFVSRKNGDLQALDNPEYHNASNGPPKAED     1200
::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::
PRGELDEEGYTMPMRDKPKQEYLNPEENPFVSRKNGDLQALDNPEYHNASNGPPKAED     433

EYVNEPLYLNTFANTLGKAEYLKNNILSMPEKAKKAFDNDYWNHSLPPRSTLQHPDYLQ     1260
::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::
EYVNEPLYLNTFANTLGKAEYLKNNILSMPEKAKKAFDNDYWNHSLPPRSTLQHPDYLQ     493

EYSTKYFYKQNGRIRPIVAENPEYLSEFSLKPGTVLPPPPYRHRNTVV      1308
::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::
EYSTKYFYKQNGRIRPIVAENPEYLSEFSLKPGTVLPPPPYRHRNTVV      541

```

Aligned 541, Matches 541, Mismatches 0, Score 130, Homology 100%

Figure 3B

[illegible]

### Figure 4


TK

HER4	680	IK-KGALNREL-ETELVEPLTFSCTAPHOALATKSTELAKVYLKAGAFOTVITUGTWVPSSTVCI PVAIKI LKSTTTO KOUWVE PDEALIMAGH	● ● ● ●
EGFR	649	V-R-T-L-Q-R.....E-L.....F-KI.....L-I.....K.....ELA.A.S.....K-I-L.....TV.....V-M	
HER2	680	Q-IR-YTN-L-Q.....AM.....M.....NK.....I-D-M.....V-R-M.S.....K-I-L.....TV.....GVS	
HER3	651	RIGH-M-Y-ENG-SI...D...-EKA-KVLA...F.....RKL.....V.....M...V.I...SI.....C...VIEDNS-ROSTDAVT-IM-AIG-L...	
HER4	778	PHLVLLGVCLSPITOLVTLPHGCLLYVHEKMDIGQOLLKNCVOIANGHTLSEDLVNRDLAANVLYKSPHRYKCTOTUGLALLDEKERTNA	▽
EGFR	748	VC.....I-TS-V-I.....F.....D...R.....D.....D.....T.Q.....K.GAE.....M	
HER2	780	YVS.....I-TS-V.....Y.....DH.R.NRGL...D...M.....S...DV.....DI...T.M	
HER3	750	A-I.....L-FGSEL.....YL-L-S...DH-RQ-RGAL-P.....GV.....Y.....KON...M...L...SO-QVA...V-D...PP-D-QILY	
HER4	878	DOORPIMHALLCTHNTKTHQSDWNSGTWELATFGKPYDGIIPFAEIPELLKBERLPQPPCTIDWTHVYKCHIDGASRPFKTELAAEPFH	
EGFR	848	E...V.....S-LH-IT.....V.....S.....AS...BST.....I.....R-II...K	
HER2	880	V.....S-LR-R.....V.....A.....A.....I.....SEC-R-R-VS...	
HER3	850	SEA-T.....S-FG-Y.....V.....AE...A-LALA-V.....A...Q.....CHIR-P.....M...T...	
HER4	978	AADPQYLVICDORHKL-PSPHDSKFTQNLLEEDELDHDAZETLYPC-AINZIPPITTAARTIDSRNEIGHPPATPHSGHGFVTRDGTAAQ	▽
EGFR	948	.....E-N.....T...N-YRA-M...MD-VV...D...I...CG-FSS...S-...	
HER2	980	.....FV...ME...LGP-A...L...T-YRS...EDO-MG-LV.....CG-FC-D-APONZVYTHGUSSTASODDL	
HER3	950	.....P.....KRESGCLA-G-EPHGLTHKK-E-VE...PEL-LDLD-EAEDWLATLGLSALFPMOTLNRPHGQSLLSPASOTYHPOQNLGDSOC	
HER4	1076	GVSVTPAFTSTIEAPVA--QDATASITDSDCHOTLANKVAPHVVEDSTORYEADPTVTAVERJPHGILDEZOTGTHMDKTRSTLNFVICHPTVS	▽
EGFR	1008	.....T-LLSLS...GN-NSTYACIDRGLQSCPIK...FL...S...GALP-D-I-----DPTL-----VZ-I-GP-----P	
HER2	1062	---GLETPSEDA-RS-L-PSE...GSDV...GQGM:ANKGLSLTHDP-FL...S...PL-S-----STD...VA-LTCS-QP...V-GTQVA-QPP	
HER3	1050	SAVGSSERCRFPVSLHPRGCLASESGHVTGSCALQCKYKCHGASRNGRPHGDSATHSGHSLLPVTFSPGSEEDWNSVTRDTLKG	▽
HER4	1174	RKNGLOALDMPETHNASNGPPKVEDTYMEPLYLNTFANTLGAETLK-----KVLNPFEDKCATONTDWNHSLPFTSLGHTLQSTRTITY	▽
EGFR	1075	K-PA-EV-Q-V...QPLH-APS-----RD-H-QDPHTAV-NP...NT-----VOTCTVSTDSF-----H-AGKSHOISLH...O-DFTS-LA-	
HER2	1151	SPNE-F-P-ANPAGATLENANTLSPPONG-VKDFP--A-GDAVEHP...TQDGAHPHNP-PAISF...LY-DQDP-E-GAPPT-----	
HER3	1150	TPSNEGTLSSVGLSYLTGTEDELECTETHRNRANSPPHPPRPSLEELQETDYNVGBDLSGLSTGSCFLNPPYINTIATTPOEDYETNRQD	
HER4	1269	KONGRIPI-VADHPEYLSFSLKPGTYLPPPYKHTVW	1308
EGFR	1158	--P--YFGS-T...A...RVAPOSSETIGA	1186
HER2	1237	---FKGPTT	1235
HER3	1250	GGPMGCDTANAGACPAEQQCTEDHRAFQCPGHQAPHVHYAALTLRLSLAETDSATONPDYHNSRLFTYKAAQRT	1313

Figure 4

(continued)

FIGURE 5







FIGURE 6




FIGURE 7




FIGURE 8



# Biological and Biochemical Properties of the MDA-MB-453-cell Differentiation Activity




FIGURE 9



FIGURE 10A





FIGURE 10B

a



H4

b




Figure 11

HER4-Ig

HER4 extracellular domain-human Ig fusion construct

**M**KPATGLWVWVSL**L**VAAGTV**Q**PSDS**Q**SVCAGTEN**K**LS**S**LS**D**LE**Q**QYRALRKYYENCEV**V**M  
**G**NLEITSIEHNRDLS**F**LR**S**VREVTGYVLVAL**N**Q**F**RYLPLE**N**LRIIRG**T**KLYEDRYALAI**F**  
**L**NYRKDGN**F**GLQELGLK**N**LTEI**L**NGGVYVD**Q**NK**F**LCYAD**T**IHWQDIVRNPWPS**N**LTLV**S**T  
**N**GSSGGRCHK**S**CTGR**C**WGPTENHC**Q**TL**T**RTVCAEQCDGR**C**YGPYSDCCHRECAGGCS**G**  
**P**KD**T**DCFAC**M**NFND**S**GACVT**Q**CP**T**FFVYN**P**TT**F**QLEHN**F**NAKYTYGAF**C**V**K**KCPH**N**FV**D**  
**S**SSCVRAC**P**SS**K**MEVEENG**I**K**M**CK**P**CTD**I**CP**K**ACDG**I**GTG**S**LS**A**QTVDSS**N**ID**K**FIN**C**T  
**K**ING**N**LI**F**LV**T**GIHGDPY**N**AI**E**AID**P**E**K**LN**V**FRTVRE**I**TG**F**LN**I**QSWPP**N**MTDFSV**F**SN**L**  
**V**TIGGRVLY**S**GL**S**LL**I**L**K**QOG**I**TS**L**Q**F**Q**S**LKEISAGNIY**I**TD**S**N**L**CY**Y**HTINW**T**TL**F**ST  
**I**NQRIVIRDN**R**KAENCTA**E**GMVC**N**HLCS**S**DGCWGPDP**Q**CL**S**CR**R**FSRGR**I**CIESCN**L**YD  
**G**EFREFENG**S**ICVECD**P**QCE**K**MEDGL**L**TCHGPGPD**N**CT**K**CS**H**FKDGPNCVE**K**CPDGL**Q**GA  
**N**S**F**IFKYADPDRECH**P**CH**P**NCT**Q**GC**N**GPT**S**HDCI**Y**YPWTGH**S**TL**P**QDPVKV**K**ALEGF**P**RL  
**V**GP**D**FFGCAEPANT**F**LD**P**EE**P**KSC**D**K**T**HTC**P**PC**P**AP**E**LLGGPSV**F**LFPP**K**PK**D**TL**M**IS**R**T  
**P**EVT**C**W**V**W**D**V**S**HED**P**EV**K**EN**T**Y**V**D**G**VE**V**Y**V**AK**T**K**P**REE**Q**Y**N**STYRVV**S**VL**T**VL**H**QDW**L**NG  
**K**EY**K**CKV**S**N**K**AL**P**API**E**KT**I**SK**A**K**Q**PRE**P**Q**V**Y**T**LP**P**SRDEL**T**KN**Q**V**S**LT**C**LV**K**G**F**Y**P**SD  
**L**AVEWESNG**Q**PEN**N**Y**K**TT**P**PVLD**S**DGS**F**FLY**S**KL**T**VD**K**SRW**Q**Q**N**V**F**SC**S**VM**H**EAL**H**N**H**Y  
**T**Q**K**SL**S**LS**P**G**K**

Bold = Signal Sequence

= Immunoglobulin domain

Lower case = HER4 ECD

Figure 12



European Patent  
Office

# PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 43 of the European Patent Convention shall be considered, for the purposes of subsequent proceedings, as the European search report

EP 93 11 8837

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cls)
P,X	WO-A-92 20798 (GENENTECH, US) 26 November 1992 * Abstract, claims *	14,15, 28,29	C12N15/12 C07K13/00 C12P21/08 C12N5/10 G01N33/68 G01N33/577 A61K39/395
P,X	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA. vol. 90 , 1 March 1993 , WASHINGTON US pages 1746 - 1750 PLOWMAN GD;CULOUSCOU JM;WHITNEY GS;GREEN JM;CARLTON GW;FOY L;NEUBAUER MG;SHOYAB M; 'Ligand-specific activation of HER4/p180erbB4, a fourth member of the epidermal growth factor receptor family.' * the whole document *	1-32	
A	SCIENCE vol. 256 , 22 May 1992 , LANCASTER, PA pages 1205 - 1210 HOLMES, W.E. ET AL.; 'Identification of heregulin, a specific activator of p185erbB2' * the whole document *	14,15	
			TECHNICAL FIELDS SEARCHED (Int.Cls)
			C12N C07K G01N
INCOMPLETE SEARCH			
<p>The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims</p> <p>Claims searched completely:</p> <p>Claims searched incompletely:</p> <p>Claims not searched:</p> <p>Reason for the limitation of the search:</p> <p>see sheet C</p>			
Place of search THE HAGUE		Date of completion of the search 9 March 1994	Examiner Nauche, S
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons A : member of the same patent family, corresponding document	

EPPO FORM 1000 (01/93)



European Patent  
Office

# PARTIAL EUROPEAN SEARCH REPORT

Application Number

EP 93 11 8837

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl. 8)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
P, X	JOURNAL OF BIOLOGICAL CHEMISTRY. vol. 268, no. 25, September 1993, BALTIMORE US pages 18407 - 18410 CULOUSCOU JM; PLOWMAN GD; CARLTON GW; GREEN JM; SHOYAB M; 'Characterization of a breast cancer cell differentiation factor that specifically activates the HER4/p180erbB4 receptor.' * page 18410, column 1, line 4 - page 18410, column 2, line 2 *	14, 15	
A	EP-A-0 444 961 (BRISTOL-MYERS SQUIBB COMPANY) 4 September 1991 * the whole document *	1-32	TECHNICAL FIELDS SEARCHED (Int. Cl. 8)
A	WO-A-90 14357 (GENENTECH, US) 29 November 1990 -----		



Europäisches  
Patentamt  
GD 1

European  
Patent Office  
DG 1

Office européen  
des brevets  
DG 1

---

SHEET C

EP 93118837.9

**Remark** : Although claim 32 is directed to a method of treatment of human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.